

# Balanced $0, \pm 1$ Matrices

## Part I: Decomposition

Michele Conforti <sup>\*</sup>  
G rard Cornu jols <sup>†</sup>  
Ajai Kapoor <sup>‡</sup>  
and Kristina Vu kovi  <sup> </sup>

revised September 2000

### Abstract

A  $0, \pm 1$  matrix is *balanced* if, in every square submatrix with two nonzero entries per row and column, the sum of the entries is a multiple of four. This paper extends the decomposition of balanced  $0, 1$  matrices obtained by Conforti, Cornu jols and Rao to the class of balanced  $0, \pm 1$  matrices. As a consequence, we obtain a polynomial time algorithm for recognizing balanced  $0, \pm 1$  matrices.

*Keywords:* balanced matrix, decomposition, recognition algorithm, 2-join, 6-join, extended star cutset

*Running head:* Decomposition of balanced  $0, \pm 1$  matrices

## 1 Introduction

A  $0, 1$  matrix is *balanced* if for every square submatrix with two ones per row and column, the number of ones is a multiple of four. This notion was introduced by Berge [1] and extended to  $0, \pm 1$  matrices by Truemper [16]. A  $0, \pm 1$  matrix is *balanced* if, in every square submatrix with two nonzero entries per row and column, the sum of the entries is a multiple of four.

This paper extends the decomposition of balanced  $0, 1$  matrices obtained by Conforti, Cornu jols and Rao [8] to the class of balanced  $0, \pm 1$  matrices. As a consequence, we obtain a polynomial time algorithm for recognizing balanced  $0, \pm 1$  matrices. The algorithm is discussed in a sequel paper.

---

<sup>\*</sup>Dipartimento di Matematica Pura ed Applicata, Universit  di Padova, Via Belzoni 7, 35131 Padova, Italy. conforti@math.unipd.it

<sup>†</sup>GSIA and Department of Mathematical Sciences, Carnegie Mellon University, Schenley Park, Pittsburgh, PA 15213. gc0v@andrew.cmu.edu

<sup>‡</sup>Dipartimento di Matematica Pura ed Applicata, Universit  di Padova, Via Belzoni 7, 35131 Padova, Italy. ajai@sfo.com

<sup> </sup>Department of Mathematics, University of Kentucky, Lexington, KY 40506. kristina@ms.uky.edu  
This work was supported in part by NSF grants DMI-9802773, DMS-9509581 and ONR grant N00014-97-1-0196.

The class of balanced  $0, \pm 1$  matrices properly includes totally unimodular  $0, \pm 1$  matrices. (A matrix is *totally unimodular* if every square submatrix has determinant equal to  $0, \pm 1$ .) The fact that every totally unimodular matrix is balanced is implied, for example, by Camion’s theorem [3] which states that a  $0, \pm 1$  matrix is totally unimodular if and only if, in every square submatrix with an *even number* of nonzero entries per row and column, the sum of the entries is a multiple of four. Therefore our work is related to Seymour’s decomposition and recognition of totally unimodular matrices [15].

In Section 3 we show that, to understand the structure of balanced  $0, \pm 1$  matrices, it is sufficient to understand the structure of their zero-nonzero pattern. Such  $0, 1$  matrices are said to be *balanceable*. Clearly balanced  $0, 1$  matrices are balanceable but the converse is not

true:  $\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$  is balanceable but not balanced. Section 5 describes the cutsets used

in our decomposition theorem and Section 6 states the theorem and outlines its proof. In Section 7, we relate our result to Seymour’s [15] decomposition theorem for totally unimodular matrices. The proofs are given in Section 8 and Section 9. The necessary definitions and notation are introduced in Section 4.

Interestingly, a number of polyhedral results known for balanced  $0, 1$  matrices and totally unimodular matrices can be generalized to balanced  $0, \pm 1$  matrices. It follows that several problems in propositional logic can be solved in polynomial time by linear programming when the underlying clauses are “balanced”. These results are reviewed in Section 2.

## 2 Bicoloring, Polyhedra and Propositional Logic

Berge [1] introduced the following notion. A  $0, 1$  matrix is *bicolorable* if its columns can be partitioned into blue and red columns in such a way that every row with two or more 1’s contains a 1 in a blue column and a 1 in a red column. This notion provides the following characterization of balanced  $0, 1$  matrices.

**Theorem 2.1** (Berge [1]) *A  $0, 1$  matrix  $A$  is balanced if and only if every submatrix of  $A$  is bicolorable.*

Ghouila-Houri [14] introduced the notion of *equitable bicoloring* for a  $0, \pm 1$  matrix  $A$  as follows. The columns of  $A$  are partitioned into blue columns and red columns in such a way that, for every row of  $A$ , the sum of the entries in the blue columns differs from the sum of the entries in the red columns by at most one.

**Theorem 2.2** (Ghouila-Houri [14]) *A  $0, \pm 1$  matrix  $A$  is totally unimodular if and only if every submatrix of  $A$  has an equitable bicoloring.*

A  $0, \pm 1$  matrix  $A$  is *bicolorable* if its columns can be partitioned into blue columns and red columns in such a way that every row with two or more nonzero entries either contains two entries of opposite sign in columns of the same color, or contains two entries of the same sign in columns of different colors. For a  $0, 1$  matrix, this definition coincides with Berge’s notion of bicoloring. Clearly, if a  $0, \pm 1$  matrix has an equitable bicoloring as defined by Ghouila-Houri, then it is bicolorable.

**Theorem 2.3** (Conforti, Cornuéjols [6]) *A  $0, \pm 1$  matrix  $A$  is balanced if and only if every submatrix of  $A$  is bicolourable.*

Balanced  $0, 1$  matrices are important in integer programming due to the fact that several polytopes, such as the set covering, packing and partitioning polytopes, only have integral extreme points when the constraint matrix is balanced. Such integrality results were first observed by Berge [2] and then expanded upon by Fulkerson, Hoffman and Oppenheim [12]. In the case of balanced  $0, \pm 1$  matrices, similar integrality results were proved by Conforti and Cornuéjols [6] for the generalized set covering, packing and partitioning polytopes.

Given a  $0, \pm 1$  matrix  $A$ , let  $n(A)$  denote the column vector whose  $i^{\text{th}}$  component is the number of  $-1$ 's in the  $i^{\text{th}}$  row of matrix  $A$ .

**Theorem 2.4** (Conforti, Cornuéjols [6]) *Let  $M$  be a  $0, \pm 1$  matrix. Then the following statements are equivalent:*

- (i)  *$M$  is balanced.*
- (ii) *For each submatrix  $A$  of  $M$ , the generalized set covering polytope  $\{x : Ax \geq \mathbf{1} - n(A), \mathbf{0} \leq x \leq \mathbf{1}\}$  is integral.*
- (iii) *For each submatrix  $A$  of  $M$ , the generalized set packing polytope  $\{x : Ax \leq \mathbf{1} - n(A), \mathbf{0} \leq x \leq \mathbf{1}\}$  is integral.*
- (iv) *For each submatrix  $A$  of  $M$ , the generalized set partitioning polytope  $\{x : Ax = \mathbf{1} - n(A), \mathbf{0} \leq x \leq \mathbf{1}\}$  is integral.*

Several problems in propositional logic can be written as generalized set covering problems. For example, the satisfiability problem in conjunctive normal form (SAT) is to find whether the formula

$$\bigwedge_{i \in S} \left( \bigvee_{j \in P_i} x_j \vee \bigvee_{j \in N_i} \neg x_j \right)$$

is true. This is the case if and only if the system of inequalities

$$\sum_{j \in P_i} x_j + \sum_{j \in N_i} (1 - x_j) \geq 1 \text{ for all } i \in S$$

has a  $0, 1$  solution vector  $x$ . This is a generalized set covering problem

$$\begin{aligned} Ax &\geq \mathbf{1} - n(A) \\ x &\in \{0, 1\}^n. \end{aligned}$$

Given a set of clauses  $\bigvee_{j \in P_i} x_j \vee \bigvee_{j \in N_i} \neg x_j$  with weights  $w_i$ , MAXSAT consists of finding a truth assignment which satisfies a maximum weight set of clauses. MAXSAT can be formulated as the integer program

$$\begin{aligned} \text{Min} \quad & \sum_{i=1}^m w_i s_i \\ & Ax + s \geq \mathbf{1} - n(A) \\ & x \in \{0, 1\}^n, s \in \{0, 1\}^m. \end{aligned}$$

Similarly, the inference problem in propositional logic can be formulated as

$$\text{min } \{cx : Ax \geq \mathbf{1} - n(A), x \in \{0, 1\}^n\}.$$

The above three problems are NP-hard in general but SAT and logical inference can be solved efficiently for Horn clauses, clauses with at most two literals and several related classes [4],[17]. MAXSAT remains NP-hard for Horn clauses with at most two literals [13]. A consequence of Theorem 2.4 is the following.

**Corollary 2.5** *SAT, MAXSAT and logical inference can be solved in polynomial time by linear programming when the corresponding  $0, \pm 1$  matrix  $A$  is balanced.*

In fact SAT and logical inference can be solved by repeated application of unit resolution when the underlying  $0, \pm 1$  matrix  $A$  is balanced [5]. These results are surveyed in [7].

### 3 Balanceable 0, 1 Matrices

In this section, we consider the following question: given a 0, 1 matrix, is it possible to turn some of the 1's into  $-1$ 's in order to obtain a balanced  $0, \pm 1$  matrix? A 0, 1 matrix for which such a signing exists is called a *balanceable* matrix.

Given a 0, 1 matrix  $A$ , the *bipartite graph representation of  $A$*  is the bipartite graph  $G = (V^r, V^c; E)$  having a node in  $V^r$  for every row of  $A$ , a node in  $V^c$  for every column of  $A$  and an edge  $ij$  joining nodes  $i \in V^r$  and  $j \in V^c$  if and only if the entry  $a_{ij}$  of  $A$  equals 1. The sets  $V^r$  and  $V^c$  are the *sides* of the bipartition. We say that  $G$  is *balanced* if  $A$  is a balanced matrix.

A *signed graph* is a graph  $G$ , together with an assignment of weights  $+1, -1$  to the edges of  $G$ . To a  $0, \pm 1$  matrix corresponds its signed bipartite graph representation. A signed bipartite graph  $G$  is *balanced* if it is the signed bipartite graph representation of a balanced  $0, \pm 1$  matrix. Thus a signed bipartite graph  $G$  is balanced if and only if, in every hole  $H$  of  $G$ , the *weight* of the hole, i.e. the sum of the weights of the edges in  $H$ , is a multiple of four. (A *hole* in a graph is a chordless cycle).

A bipartite graph  $G$  is *balanceable* if there exists a signing of its edges so that the resulting signed graph is balanced.

**Remark 3.1** *Since cuts and cycles of a connected graph have even intersection, it follows that, if a connected signed bipartite graph  $G$  is balanced, then the signed bipartite graph  $G'$ , obtained by switching signs on the edges of a cut, is also balanced.*

For every edge  $uv$  of a spanning tree, there is a cut containing  $uv$  and no other edge of the tree (such cuts are known as *fundamental cuts*). Thus, if  $G$  is a connected balanceable bipartite graph, the edges of a spanning tree can be signed arbitrarily and then the remaining edges can still be signed so that  $G$  is a balanced signed bipartite graph. This was already observed by Camion [3] in the context of 0, 1 matrices that can be signed to be totally unimodular. So Remark 3.1 implies that a bipartite graph  $G$  is balanceable if and only if the following signing algorithm produces a balanced signed bipartite graph:

#### Signing Algorithm

*Choose a spanning forest of  $G$ , sign its edges arbitrarily and recursively choose an edge  $uv$ , which closes a hole  $H$  of  $G$  with the previously chosen edges, and sign  $uv$  so that the sum of the weights of the edges in  $H$  is a multiple of four.*




Figure 1: 3-path configuration and odd wheel

Note that, in the signing algorithm, the edge  $uv$  can be chosen to close the smallest length hole with the previously chosen edges. Such a hole  $H$  is also a hole in  $G$ .

It follows from this signing algorithm that, up to the signing of a spanning forest, a balanceable bipartite graph has only one signing that makes it balanced. Consequently, the problem of recognizing whether a bipartite graph is balanceable is equivalent to the problem of recognizing whether a signed bipartite graph is balanced.

Let  $G$  be a bipartite graph. Let  $u, v$  be two nonadjacent nodes in opposite sides of the bipartition. A *3-path configuration connecting  $u$  and  $v$* , denoted by  $3PC(u, v)$ , is defined by three chordless paths  $P_1, P_2, P_3$  with endnodes  $u$  and  $v$ , such that the node set  $V(P_i) \cup V(P_j)$  induces a hole for  $i \neq j$  and  $i, j \in \{1, 2, 3\}$ . In particular, none of the three paths is an edge. A 3-path configuration is shown in Figure 1. (In all figures black nodes and white nodes are nodes on opposite sides of the bipartition. A solid line denotes an edge, a dashed one a path that is not an edge). Since paths  $P_1, P_2, P_3$  of a 3-path configuration are of length one or three modulo four, the sum of the weights of the edges in each path is also one or three modulo four. It follows that two of the three paths induce a hole of weight two modulo four. So a bipartite graph which contains a 3-path configuration as an induced subgraph is not balanceable.

A *wheel*, denoted by  $(H, x)$ , is defined by a hole  $H$  and a node  $x \notin V(H)$  having at least three neighbors in  $H$ , say  $x_1, x_2, \dots, x_n$ . If  $n$  is even, the wheel is an *even wheel*, otherwise it is an *odd wheel* (for example see Figure 1). An edge  $xx_i$  is a *spoke*. A subpath of  $H$  connecting  $x_i$  and  $x_j$  is called a *sector* if it contains no intermediate node  $x_l, 1 \leq l \leq n$ . Consider a wheel which is signed to be balanced. By Remark 3.1, all spokes of the wheel can be assumed to have weight 1. This implies that the sum of the weights of the edges in each sector is two modulo four. Hence the wheel must be an even wheel else the hole  $H$  has weight two modulo four.

So, balanceable bipartite graphs contain neither odd wheels nor 3-path configurations as induced subgraphs. This fact is used extensively in our proofs in this paper. The following theorem of Truemper [16] states that the converse is also true.

**Theorem 3.2** (Truemper [16]) *A bipartite graph is balanceable if and only if it does not contain an odd wheel nor a 3-path configuration as an induced subgraph.*

## 4 Additional Definitions and Notation

Let  $G$  be a bipartite graph where the two sides of the bipartition are  $V^r$  and  $V^c$ .  $G$  contains a graph  $\Sigma$  if  $\Sigma$  is an induced subgraph of  $G$ .  $N(v)$  refers to the set of nodes adjacent to node  $v$ . A node  $v \notin V(\Sigma)$  is *strongly adjacent* to  $\Sigma$  if  $|N(v) \cap V(\Sigma)| \geq 2$ . A node  $v$  is a *twin* of a node  $x \in V(\Sigma)$  with respect to  $\Sigma$  if  $N(v) \cap V(\Sigma) = N(x) \cap V(\Sigma)$ .

A *path*  $P$  is a sequence of distinct nodes  $x_1, x_2, \dots, x_n$ ,  $n \geq 1$ , such that  $x_i x_{i+1}$  is an edge, for all  $1 \leq i < n$ . Let  $x_i$  and  $x_l$  be two nodes of  $P$ , where  $l \geq i$ . The path  $x_i, x_{i+1}, \dots, x_l$  is called the  $x_i x_l$ -subpath of  $P$  and is denoted by  $P_{x_i x_l}$ . We write  $P = x_1, \dots, x_{i-1}, P_{x_i x_l}, x_{l+1}, \dots, x_n$  or  $P = x_1, \dots, x_i, P_{x_i x_l}, x_l, \dots, x_n$ . A cycle  $C$  is a sequence of nodes  $x_1, x_2, \dots, x_n, x_1$ ,  $n \geq 3$ , such that the nodes  $x_1, x_2, \dots, x_n$  form a path and  $x_1 x_n$  is an edge. The node set of a path or a cycle  $Q$  is denoted by  $V(Q)$ .

Let  $A, B, C$  be three disjoint node sets such that no node of  $A$  is adjacent to a node of  $B$ . A path  $P = x_1, x_2, \dots, x_n$  *connects*  $A$  and  $B$  if one of the two endnodes of  $P$  is adjacent to at least one node in  $A$  and the other is adjacent to at least one node in  $B$ . The path  $P$  is a *direct connection between*  $A$  and  $B$  if, in the subgraph induced by the node set  $V(P) \cup A \cup B$ , no path connecting  $A$  and  $B$  is shorter than  $P$ . A direct connection  $P$  between  $A$  and  $B$  *avoids*  $C$  if  $V(P) \cap C = \emptyset$ . The direct connection  $P$  is said to be *from*  $A$  *to*  $B$  if  $x_1$  is adjacent to some node in  $A$  and  $x_n$  to some node in  $B$ .

## 5 Cutsets

In this section we introduce the operations needed for our decomposition result. A set  $S$  of nodes (edges) of a connected graph  $G$  is a *node cutset* (an *edge cutset* respectively) if the subgraph  $G \setminus S$ , obtained from  $G$  by removing the nodes (edges) in  $S$ , is disconnected.

### Extended Star Cutsets

A *biclique* is a complete bipartite graph  $K_{AB}$  where the two sides of the bipartition  $A$  and  $B$  are both nonempty.

In a connected bipartite graph  $G$ , an *extended star*  $(x; T; A; R)$  is defined by disjoint subsets  $T, A, R$  of  $V(G)$  and a node  $x \in T$  such that

- (i)  $A \cup R \subseteq N(x)$ ,
- (ii) the node set  $T \cup A$  induces a biclique (with node set  $T$  on one side of the bipartition and node set  $A$  on the other),
- (iii) if  $|T| \geq 2$ , then  $|A| \geq 2$ .

This concept was introduced in [8]. An *extended star cutset* is an extended star  $(x; T; A; R)$  where  $T \cup A \cup R$  is a node cutset. When  $R = \emptyset$  the extended star is a biclique, and the cutset is called a *biclique cutset*.

## Joins

Let  $G$  be a connected bipartite graph containing a biclique  $K_{A_1A_2}$  with the property that its edge set  $E(K_{A_1A_2})$  is a cutset of  $G$  and no connected component of  $G' = G \setminus E(K_{A_1A_2})$  contains both a node of  $A_1$  and a node of  $A_2$ . For  $i = 1, 2$ , let  $G'_i$  be the union of the components of  $G'$  containing a node of  $A_i$ . The edge set  $E(K_{A_1A_2})$  is a *1-join* if the graphs  $G'_1$  and  $G'_2$  each contains at least two nodes. This concept was introduced by Cunningham and Edmonds [11].

Let  $G$  be a connected bipartite graph with more than four nodes, containing bicliques  $K_{A_1A_2}$  and  $K_{B_1B_2}$ , where  $A_1, A_2, B_1, B_2$  are disjoint nonempty node sets. The edge set  $E(K_{A_1A_2}) \cup E(K_{B_1B_2})$  is a *2-join* if it satisfies the following properties:

- (i) The graph  $G' = G \setminus (E(K_{A_1A_2}) \cup E(K_{B_1B_2}))$  is disconnected.
- (ii) Every connected component of  $G'$  has a nonempty intersection with exactly two of the sets  $A_1, A_2, B_1, B_2$  and these two sets are either  $A_1$  and  $B_1$  or  $A_2$  and  $B_2$ . For  $i = 1, 2$ , let  $G'_i$  be the subgraph of  $G'$  containing all its connected components that have nonempty intersection with  $A_i$  and  $B_i$ .
- (iii) If  $|A_1| = |B_1| = 1$ , then  $G'_1$  is not a chordless path or  $A_2 \cup B_2$  induces a biclique. If  $|A_2| = |B_2| = 1$ , then  $G'_2$  is not a chordless path or  $A_1 \cup B_1$  induces a biclique.

This concept was introduced by Cornuéjols and Cunningham [10] and was extensively used in [8]. In the present paper, 2-joins are needed in the statement of the main theorem, which builds on the work of [8], but do not occur in the proofs.

In a connected bipartite graph  $G$ , let  $A_i, i = 1, \dots, 6$  be disjoint, nonempty node sets such that, for each  $i$ , every node in  $A_i$  is adjacent to every node in  $A_{i-1} \cup A_{i+1}$  (indices are taken modulo 6), and these are the only edges in the subgraph  $A$  induced by the node set  $\cup_{i=1}^6 A_i$ . (Note that for convenience of notation the modulo 6 function is assumed to return values between 1 and 6, instead of the usual 0 to 5). The edge set  $E(A)$  is a *6-join* if

- (i) The graph  $G' = G \setminus E(A)$  is disconnected.
- (ii) The nodes of  $G$  can be partitioned into  $V_T$  and  $V_B$  so that  $A_1 \cup A_3 \cup A_5 \subseteq V_T, A_2 \cup V_4 \cup V_6 \subseteq V_B$  and the only adjacencies between  $V_T$  and  $V_B$  are the edges of  $E(A)$ .
- (iii)  $|V_T| \geq 4$  and  $|V_B| \geq 4$ .

When the graph  $G$  comprises more than one connected component, we say that  $G$  has a 1-join, a 2-join, a 6-join or an extended star cutset if at least one of its connected components does.

## 6 The Main Theorem

A bipartite graph is *strongly balanceable* if it is balanceable and contains no cycle with exactly one chord. Strongly balanceable bipartite graphs can be recognized in polynomial time [9].  $R_{10}$  is the balanceable bipartite graph defined by the cycle  $x_1, \dots, x_{10}, x_1$  of length 10 with




Figure 2: A 6-join




Figure 3:  $R_{10}$

chords  $x_i x_{i+5}$ ,  $1 \leq i \leq 5$  (see Figure 3). For example, a proper signing of  $R_{10}$  is to assign weight  $+1$  to the edges of the cycle  $x_1, \dots, x_{10}, x_1$  and  $-1$  to the chords.

We can now state the decomposition theorem for balanceable bipartite graphs:

**Theorem 6.1** *A balanceable bipartite graph that is not strongly balanceable is either  $R_{10}$  or contains a 2-join, a 6-join or an extended star cutset.*

The key idea in the proof of Theorem 6.1 is that if a balanceable bipartite graph  $G$  is not strongly balanceable, then one of the three following cases occurs: (i) the graph  $G$  contains  $R_{10}$  as an induced subgraph, or (ii) it contains a certain induced subgraph which forces a 6-join or an extended star cutset of  $G$ , or (iii) an earlier result of Conforti, Cornuéjols and Rao [8] applies.

### Connected 6-Holes

A *triad* is a bipartite graph consisting of three internally node-disjoint paths  $t, \dots, u$ ;  $t, \dots, v$  and  $t, \dots, w$ , where  $t, u, v, w$  are distinct nodes and belong to the same side of the bipartition. Furthermore, the graph induced by the nodes of the triad contains no other edges than those of the three paths. Nodes  $u, v$  and  $w$  are called the *attachments* and  $t$  is called the *meet* of the triad.

A *fan* consists of a chordless path  $P = x, \dots, y$  together with a node  $z$  not in  $P$  adjacent to a positive even number of nodes in  $P$ , where  $x, y$  and  $z$  belong to the same side of the bipartition and are called the *attachments* of the fan. Node  $z$  is the *center* of the fan and the edges connecting  $z$  to  $P$  are the *spokes*.

A *connected 6-hole*  $\Sigma$  is a bipartite graph induced by two disjoint node sets  $T(\Sigma)$  and  $B(\Sigma)$  such that each induces either a triad or a fan, the attachments of  $B(\Sigma)$  and  $T(\Sigma)$  induce a 6-hole and there are no other adjacencies between the nodes of  $T(\Sigma)$  and  $B(\Sigma)$ .  $T(\Sigma)$  and  $B(\Sigma)$  are the *sides* of  $\Sigma$ ,  $T(\Sigma)$  is the *top* and  $B(\Sigma)$  the *bottom*.

In this paper we will prove the following two theorems.

**Theorem 6.2** *A balanceable bipartite graph that contains  $R_{10}$  as a proper induced subgraph has a biclique cutset.*

**Theorem 6.3** *A balanceable bipartite graph that contains a connected 6-hole as an induced subgraph has an extended star cutset or a 6-join.*

Now Theorem 6.1 follows from Theorems 6.2, 6.3 and the following result.

**Theorem 6.4** [8] *A balanceable bipartite graph not containing  $R_{10}$  or a connected 6-hole as induced subgraphs either is strongly balanceable or contains a 2-join or an extended star cutset.*

A signed bipartite graph is *strongly balanced* if it is balanced and contains no cycle with exactly one chord. A corollary of Theorem 6.1 and of the signing algorithm is the following result.

**Theorem 6.5** *A signed bipartite graph that is balanced but not strongly balanced is either  $R_{10}$  with proper signing or it contains a 2-join, a 6-join or an extended star cutset.*

**Conjecture 6.6** *If a  $0, \pm 1$  matrix is balanced but not totally unimodular, then the underlying signed bipartite graph contains an extended star cutset.*

The restriction of this conjecture to  $0, 1$  matrices is true: a proof can be found in [8].

## 7 Connection with Seymour's Decomposition of Totally Unimodular Matrices

Seymour [15] discovered a decomposition theorem for  $0,1$  matrices that can be signed to be totally unimodular. The decompositions involved in his theorem are 1-separations, 2-separations and 3-separations. A matrix  $B$  has a  $k$ -separation if its rows and columns can be partitioned so that, after permutation of rows and columns,

$$B = \begin{pmatrix} A^1 & D^2 \\ D^1 & A^2 \end{pmatrix}$$

where  $r(D^1) + r(D^2) = k - 1$  and the number of rows plus number of columns of  $A^i$  is at least  $k$ , for  $i = 1, 2$  (here  $r(C)$  denotes the  $\text{GF}(2)$ -rank of the  $0,1$  matrix  $C$ ).

For a 1-separation,  $r(D^1) + r(D^2) = 0$ . Thus both  $D^1$  and  $D^2$  are identically zero. The bipartite graph corresponding to the matrix  $B$  is disconnected.

For the 2-separation,  $r(D^1) + r(D^2) = 1$ , thus w.l.o.g.  $D^2$  has rank zero and is identically zero. Since  $r(D^1) = 1$ , after permutation of rows and columns,  $D^1 = \begin{pmatrix} \mathbf{0} & E \\ \mathbf{0} & \mathbf{0} \end{pmatrix}$ , where  $E$  is a matrix all of whose entries are 1. The 2-separation in the bipartite graph representation of  $B$  corresponds to a 1-join.

For the 3-separation,  $r(D^1) + r(D^2) = 2$ . If both  $D^1$  and  $D^2$  have rank 1 then, after permutation of rows and columns,

$$D^1 = \begin{pmatrix} \mathbf{0} & E^1 \\ \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad D^2 = \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ E^2 & \mathbf{0} \end{pmatrix}$$

where  $E^1$  and  $E^2$  are matrices whose entries are all 1. This 3-separation in the bipartite graph representation of  $B$  corresponds to a 2-join.

When  $r(D^1) = 2$  or  $r(D^2) = 2$ , it can be shown that the resulting 3-separation corresponds to a 2-join, a 6-join or to one of two other decompositions which each contain an extended star cutset.

In order to prove his decomposition theorem, Seymour used matroid theory. A matroid is *regular* if it is binary and its partial representations can be signed to be totally unimodular (see [18] for relevant definitions in matroid theory). The elementary families in Seymour's decomposition theorem consist of graphic matroids, cographic matroids and a 10-element matroid called  $\mathcal{R}_{10}$ .  $\mathcal{R}_{10}$  has exactly two partial representations

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 1 \end{pmatrix}$$

The bipartite graph representations are shown in Figure 4.

**Theorem 7.1** (Seymour [15]) *A regular matroid is either graphic, cographic, the 10-element matroid  $\mathcal{R}_{10}$ , or it contains a 1-, 2- or 3-separation.*




Figure 4: Representations of  $\mathcal{R}_{10}$

In order to prove Theorem 7.1, Seymour first showed that a regular matroid which is not graphic or cographic either contains a 1- or 2-separation or contains an  $\mathcal{R}_{10}$  or an  $\mathcal{R}_{12}$  minor, where  $\mathcal{R}_{12}$  is a 12-element matroid having the following matrix as one of its partial representations.

$$\begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix}$$

Note that the bipartite graph representation of this matrix is a connected 6-hole where both sides are fans. So, this first part in Seymour's proof has some similarity with Theorem 6.4 stated above for balanceable bipartite graphs.

Then Seymour showed that, if a regular matroid contains an  $\mathcal{R}_{10}$  minor, either it is  $\mathcal{R}_{10}$  itself or it contains a 1-separation or a 2-separation. This is similar to Theorem 6.2.

Seymour completed his proof by showing that, for a regular matroid which contains an  $\mathcal{R}_{12}$  minor, the 3-separation of  $\mathcal{R}_{12}$  induces a 3-separation for the matroid. We show that for a balanceable bipartite graph, which contains a connected 6-hole as an induced subgraph, either the 6-join of the connected 6-hole induces a 6-join of the whole graph or there is an extended star cutset (Theorem 6.3).

Our proof differs significantly from Seymour's for the following reason: a regular matroid may have a large number of partial representations which lead to nonisomorphic bipartite graphs. This is the case for  $\mathcal{R}_{12}$ . All these partial representations are related through pivoting. In the case of  $0,1$  balanceable matrices there is no underlying matroid, so pivoting cannot help reduce the number of cases. Since our proof is broken down differently from Seymour's, we do not consider all these cases explicitly either.

## 8 Splitter Theorem for $R_{10}$

An *extended*  $R_{10}$  is a bipartite graph induced by ten nonempty pairwise disjoint node sets  $T_1, \dots, T_{10}$  such that for every  $1 \leq i \leq 10$ , the node sets  $T_i \cup T_{i-1}$ ,  $T_i \cup T_{i+1}$  and  $T_i \cup T_{i+5}$  all induce bicliques and these are the only edges in the graph. Throughout this section, all the indices are taken modulo 10.

We consider a balanceable bipartite graph  $G$  which contains a node induced subgraph  $R$  isomorphic to  $R_{10}$ . We denote its node set by  $\{1, \dots, 10\}$  and for each  $i = 1, \dots, 10$ , node  $i$  is adjacent to nodes  $i-1, i+1$  and  $i+5$ .

In this section we give a proof of Theorem 6.2. The first step in the proof of the theorem is to study the structure of the strongly adjacent nodes to  $R$ .

**Theorem 8.1** *Let  $R$  be an  $R_{10}$  of  $G$ . If  $w$  is a strongly adjacent node to  $R$ , then  $w$  is a twin of a node in  $V(R)$  with respect to  $R$ .*

*Proof:* First, assume that  $w$  has exactly two neighbors in  $R$ . If the neighbors of  $w$  in  $R$  are nodes 1 and 3, the hole  $w, 1, 6, 7, 8, 3, w$  induces an odd wheel with center 2. If the neighbors of  $w$  in  $R$  are nodes 1 and 5, the hole  $w, 1, 2, 7, 8, 9, 4, 5, w$  is an odd wheel with center 10. The other cases where  $w$  has two neighbors in  $R$  are isomorphic.

We now assume that node  $w$  is adjacent to at least three nodes in  $R$ . If node  $w$  is adjacent to nodes  $i, i+2, i+4$ , then there exists an odd wheel  $i, i+1, i+2, i+3, i+4, i+5, i$  with center  $w$ . So  $w$  is adjacent to exactly three nodes  $i, i+2, i+6$ , showing that  $w$  is a twin of  $i+1$ .  $\square$

**Definition 8.2** *Let  $R$  be an  $R_{10}$  of  $G$ . For  $1 \leq i \leq 10$ , let  $T_i(R)$  be the set of nodes comprising node  $i$  in  $R$  and all the twins of node  $i$  with respect to  $R$ . Let  $R^*$  be the graph induced by the node set  $\cup_{i=1}^{10} T_i(R)$ .*

**Lemma 8.3**  *$R^*$  is an extended  $R_{10}$ .*

*Proof:* Let  $u \in T_i(R)$  and  $v \in T_j(R)$ , where  $1 \leq i, j \leq 10$ . Let  $R'$  be the  $R_{10}$  obtained from  $R$  by substituting node  $u$  for node  $i$ . Now by Theorem 8.1, node  $v$  is twin of node  $j$  in  $R'$ . Hence nodes  $u$  and  $v$  are adjacent if and only if nodes  $i$  and  $j$  are adjacent.  $\square$

**Theorem 8.4**  *$R^*$  satisfies the following two properties:*

- (i) *If node  $w$  is strongly adjacent to  $R^*$  then for some  $1 \leq i \leq 10$ ,  $N(w) \cap V(R^*) \subseteq T_i(R)$ .*
- (ii) *If  $R'$  is an  $R_{10}$  induced by the node set  $\{x_1, \dots, x_{10}\}$  where  $x_i \in T_i(R)$  for  $1 \leq i \leq 10$ , then  $T_i(R') = T_i(R)$ .*

*Proof:* To prove (i), assume that  $w$  is adjacent to  $w_i \in T_i(R)$  and  $w_j \in T_j(R)$ ,  $i \neq j$ . Let  $R_{w_i w_j}$  be an  $R_{10}$  obtained from  $R$  by replacing node  $i$  with  $w_i$  and node  $j$  with  $w_j$ . Node  $w$  is now strongly adjacent to  $R_{w_i w_j}$ , so by Theorem 8.1, node  $w$  is a twin of a node in  $R_{w_i w_j}$ . Hence  $w$  is adjacent to a node  $k$  of  $R$ . Let  $R_{w_i}$  be an  $R_{10}$  obtained from  $R$  by replacing node  $i$  by  $w_i$ . Since  $w$  is adjacent to  $k$  and  $w_i$ , it is strongly adjacent to  $R_{w_i}$ , hence, by Theorem 8.1,  $w$  is adjacent to a node  $l \neq k$  of  $R$ . Now  $w$  is a strongly adjacent node of  $R$  and, by Theorem 8.1, must be a twin of a node of  $R$ . Hence  $w \in V(R^*)$ , which contradicts our choice of  $w$ .

To prove (ii), note that Lemma 8.3 implies  $T_i(R) \subseteq T_i(R')$ , so it is enough to show that  $T_i(R') \subseteq T_i(R)$ . Let  $u \in T_i(R')$  and suppose that  $u \notin T_i(R)$ . Then node  $u$  is strongly adjacent to  $R^*$  and by (i) we have a contradiction.  $\square$

**Remark 8.5** *Considering Theorem 8.4, we can simplify the notation by replacing  $T_i(R)$  by  $T_i$ .*

**Definition 8.6** *For  $1 \leq i \leq 10$ , let  $K_i$  be the complete bipartite graph induced by the node set  $T_{i-1} \cup T_i \cup T_{i+1} \cup T_{i+5}$ .*

We now study the structure of paths between the nodes of  $R^*$ .

**Lemma 8.7** *If  $P = x_1, \dots, x_n$  is a direct connection from  $T_i$  to  $V(R^*) \setminus T_i$  in  $G \setminus E(K_i)$ , then the neighbors of  $x_n$  in  $R^*$  belong to a unique set  $T_j$ , where  $j = i - 1, i + 1$  or  $i + 5$ .*

*Proof:* Assume w.l.o.g. that  $x_1$  is adjacent to node  $i$ . By Theorem 8.4 (i),  $n > 1$  and node  $x_n$  has neighbors in exactly one  $T_j$ . Assume that for some  $j \notin \{i - 1, i + 1, i + 5\}$ ,  $x_n$  is adjacent to a node  $v_j \in T_j$ .

If  $j = i + 2$  then the hole  $i, x_1, P, x_n, v_{i+2}, i + 7, i + 6, i + 5, i$  induces an odd wheel with center  $i + 1$ . If  $j = i + 3$  then the paths  $P_1 = i, x_1, P, x_n, v_{i+3}; P_2 = i, i + 1, i + 2, v_{i+3}$  and  $P_3 = i, i - 1, i + 4, v_{i+3}$  induce a  $3PC(i, v_{i+3})$ . If  $j = i + 4$  then the hole  $i, x_1, P, x_n, v_{i+4}, i + 3, i + 8, i + 7, i + 6, i + 1, i$  induces an odd wheel with center  $i + 2$ . This completes the proof since the remaining cases are isomorphic to the above three.  $\square$

**Lemma 8.8** *There cannot exist a path  $P = x_1, \dots, x_n$  with nodes belonging to  $V(G) \setminus V(R^*)$  such that  $x_1$  is adjacent to a node  $v_i \in T_i$  and  $x_n$  is adjacent to a node  $v_j \in T_j$ , where  $i \neq j$  and  $v_i$  and  $v_j$  are not adjacent.*

*Proof:* Let  $P$  be a shortest path contradicting the lemma. Hence  $P$  does not contain an intermediate node adjacent to a node in  $T_i \cup T_j$ . By Theorem 8.1,  $n > 1$ . If no node  $x_l$  of  $P$ ,  $2 \leq l \leq n - 1$ , is adjacent to a node in  $V(R^*)$  then  $P$  is a direct connection from  $T_i$  to  $V(R^*) \setminus T_i$  in  $G \setminus E(K_i)$  contradicting Lemma 8.7.

Let  $w \in T_k$ ,  $k \neq i, j$ , be adjacent to a node of  $P$ . By minimality of  $P$ ,  $w$  is adjacent to  $v_i$  and  $v_j$  and no node of  $V(P) \setminus \{x_1, x_n\}$  is adjacent to a node of  $V(R^*) \setminus T_k$ . By symmetry, there are two cases to consider:  $k, j$  are either  $i + 1, i + 2$  or  $i + 1, i + 6$ . In the first case, let  $H_1 = v_i, P, v_{i+2}, i + 3, i + 4, i - 1, v_i$  and  $H_2 = v_i, P, v_{i+2}, i + 7, i + 6, i + 5, v_i$ . Now either  $H_1$  or  $H_2$  induces an odd wheel with center  $w$  depending on the number of neighbors of  $w$  in  $P$ . In the second case, the hole  $v_i, P, v_{i+6}, i + 7, i + 2, i + 3, i + 4, i - 1, v_i$  induces an odd wheel with center  $i + 5$ .  $\square$

*Proof of Theorem 6.2:* Let  $G$  be a balanceable bipartite graph. Let  $R$  be an  $R_{10}$  of  $G$ . By Lemma 8.3,  $R^*$  is an extended  $R_{10}$ . Assume that  $V(G) \neq V(R^*)$ . Let  $w$  be a node in  $V(G) \setminus V(R^*)$  adjacent to a node in  $T_i$ . If the biclique  $K_i$  is not a cutset of  $G$ , separating  $w$  from  $V(R^*)$ , then a path contradicting Lemma 8.8 exists. Hence  $V(G) = V(R^*)$ . If  $G$  is not  $R_{10}$ , then at least one of the node sets  $T_i(R)$  has cardinality greater than one. W.l.o.g. let  $u$  and  $v$  be two nodes in  $T_1(R)$ . Now  $\{u\} \cup N(u)$  is a biclique cutset separating  $v$  from the rest of the graph.  $\square$

## 9 Connected 6-Hole

Let  $\Sigma$  be a connected 6-hole induced by  $T(\Sigma)$  and  $B(\Sigma)$  in a balanceable bipartite graph  $G$ . In this section, we prove that either  $G$  contains an extended star cutset or it has a 6-join which separates the top and the bottom of  $\Sigma$  (Theorem 6.3).

We denote by  $H = h_1, h_2, h_3, h_4, h_5, h_6, h_1$  the 6-hole of  $\Sigma$  and we assume that  $h_1, h_3, h_5 \in T(\Sigma)$  and  $h_2, h_4, h_6 \in B(\Sigma)$ . We also assume  $h_1, h_3, h_5 \in V^c$  and  $h_2, h_4, h_6 \in V^r$ . Throughout the remainder of the paper indices referring to the hole will be taken modulo 6. If  $T(\Sigma)$  is a triad, then the three paths defining it are denoted by  $P_1, P_3$  and  $P_5$  with endnodes  $h_1, h_3$  and  $h_5$  respectively and the meet is denoted by  $t$ .

The idea of the proof is to extend the 6-join of  $\Sigma$  into a 6-join of  $G$ . Namely, we aim to find node sets  $H_1, H_2, \dots, H_6$  such that  $h_i \in H_i$ , for  $1 \leq i \leq 6$ , and  $E(\cup_{i=1}^6 H_i)$  is a 6-join for  $G$  separating  $T(\Sigma)$  from  $B(\Sigma)$ . If this is not possible, we detect an extended star cutset in  $G$ .

**Remark 9.1** *Let  $h_i$  and  $h_j$  be two distinct attachments of a side  $X$  of  $\Sigma$ . There is a unique chordless path in  $X$ , connecting  $h_i$  and  $h_j$ . This path is denoted by  $P_{ij}$ . Also any pair of nodes in  $V(\Sigma)$  are contained in a hole of  $\Sigma$ .*

**Definition 9.2** *A tripod with attachments  $x, y, z$  is either a fan where we allow the center to have any positive number (even or odd) of neighbors in the path  $P$ , or a triad where the meet is not adjacent to any of the attachments but is not restricted to be in the same side of the bipartition as the attachments.*

**Lemma 9.3** *Let  $G$  be a bipartite graph and let  $x, y, z$  be distinct nodes in the same side of the bipartition such that both  $G$  and  $G \setminus \{x, y, z\}$  are connected. Then  $G$  contains a tripod with attachments  $x, y$  and  $z$ .*

*Proof:* Let  $G'$  be a minimal subgraph of  $G$  such that  $x, y, z$  are in  $G'$  and both  $G', G' \setminus \{x, y, z\}$  are connected. We show that  $G'$  is a tripod with attachments  $x, y, z$ .

Let  $P_{xy} = x, y_1, \dots, y_m, y$  be a shortest  $xy$ -path in  $G' \setminus \{z\}$ ,  $P_{xz}$  and  $P_{yz}$  similarly defined. Assume w.l.o.g. that  $P_{xy}$  is not shorter than any of the other two. If  $P_{xy}$  contains a neighbor of  $z$  then  $V(G') = V(P_{xy}) \cup \{z\}$  and  $G'$  is a tripod. Otherwise let  $P_z = x_1, \dots, x_n$ , be a direct connection in  $G'$  from  $z$  to  $V(P_{xy}) \setminus \{x, y\}$ . By the minimality of  $G'$ ,  $V(G') = V(P_{xy}) \cup V(P_z)$  and  $x_n$  has a unique neighbor, say  $x_{n+1}$ , in  $P_{xy}$ . If  $x$  has a neighbor in  $P_z$ , by the minimality of  $G'$  and the fact that  $V(G') = V(P_{xy}) \cup V(P_z)$ ,  $y$  has no neighbor in  $P_z$  and  $x_{n+1}$  is adjacent to  $x$ . Now this contradicts our choice of  $P_{xy}$ .

By symmetry, neither  $x$  nor  $y$  have neighbors in  $P_z$  and if  $x_{n+1}$  is adjacent to  $x$  or  $y$  our choice of  $P_{xy}$  is contradicted, so  $G'$  is a tripod.  $\square$

**Lemma 9.4** *In a balanceable bipartite graph  $G$ , let  $T$  and  $B$  be node disjoint tripods with attachments  $h_1, h_3, h_5$  and  $h_2, h_4, h_6$ . If  $h_1, h_2, h_3, h_4, h_5, h_6, h_1$  is a 6-hole of  $G$  and no other adjacency exists between  $T$  and  $B$ , then  $T$  is a fan or a triad and so is  $B$ . Therefore  $T$  and  $B$  are the top and bottom of a connected 6-hole.*

*Proof:* Let  $\Sigma$  be the graph induced by  $V(T) \cup V(B)$ . Let  $P_{13}$  be a shortest  $h_1h_3$ -path in  $T \setminus \{h_5\}$ . If  $P_{13}$  contains neighbors of  $h_5$  and  $T$  is not a fan, then  $\Sigma$  contains an odd wheel. So by symmetry we can assume that  $T$  contains three chordless paths  $t, \dots, h_1, t, \dots, h_3$  and  $t, \dots, h_5$  and  $t$  not adjacent to any of the nodes  $h_1, h_3$  and  $h_5$ . If  $t$  and  $h_1$  are on opposite sides of the bipartition,  $\Sigma$  contains a  $3PC(t, h_1)$ . Therefore  $T$  is a triad. Similarly,  $B$  is a fan or a triad.  $\square$

## 9.1 Strongly Adjacent Nodes and Direct Connections

**Theorem 9.5** *Let  $\Sigma$  be a connected 6-hole in a balanceable bipartite graph  $G$ . Let  $P = x_1, \dots, x_n$  (we allow  $n = 1$ ) be a direct connection between  $T(\Sigma)$  and  $B(\Sigma)$  in  $G \setminus E(H)$  such that either  $x_1$  has a neighbor in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  or  $x_n$  has a neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$  or both. Then either  $x_1$  has exactly two neighbors in  $\{h_1, h_3, h_5\}$  and no other neighbor in  $T(\Sigma)$  or  $x_n$  has exactly two neighbors in  $\{h_2, h_4, h_6\}$  and no other neighbor in  $B(\Sigma)$ .*

To prove the above theorem, we use the following result about the structure of strongly adjacent nodes to an even wheel. Two sectors of a wheel  $(W, v)$  are *adjacent* if they have a common endnode. A *bicoloring* of  $(W, v)$  is an assignment of two colors to the intermediate nodes of its sectors so that the nodes in the same sector have the same color and nodes of adjacent sectors have distinct colors. The neighbors of  $v$  are left unpainted. Note that a wheel is bicolorable if and only if it is even.

**Lemma 9.6** *Let  $(W, v)$ ,  $v \in V^r$ , be a bicolored wheel in a balanceable bipartite graph, and let  $u \in V^c \setminus N(v)$  be a node with neighbors in at least two distinct sectors of the wheel  $(W, v)$ . Then  $u$  satisfies one of the following properties:*

**Type a** *Node  $u$  has exactly two neighbors in  $W$  and these neighbors belong to two distinct sectors having the same color.*

**Type b** *There exists one sector, say  $S_j$  with endnodes  $v_i$  and  $v_k$ , such that  $u$  has a positive even number of neighbors in  $S_j$  and has exactly two neighbors in  $V(W) \setminus V(S_j)$ , adjacent to  $v_i$  and  $v_k$  respectively.*

*Proof:* Assume first that  $u$  has neighbors in at least three different sectors, say  $S_i, S_j, S_k$ . If none of these sectors is adjacent to both of the other two, then there exist three unpainted nodes  $v_i, v_j, v_k$ , such that  $v_i \in V(S_i) \setminus (V(S_j) \cup V(S_k))$ ,  $v_j \in V(S_j) \setminus (V(S_i) \cup V(S_k))$ ,  $v_k \in V(S_k) \setminus (V(S_i) \cup V(S_j))$ . This implies the existence of a  $3PC(u, v)$ , where each of the nodes  $v_i, v_j, v_k$  belongs to a distinct path of the 3-path configuration. So  $u$  has neighbors in exactly three sectors and one of them is adjacent to the other two, say  $S_j$  is adjacent to both  $S_i$  and  $S_k$ . Let  $v_i$  be the unpainted node in  $V(S_i) \cap V(S_j)$  and  $v_k$  the unpainted node in  $V(S_j) \cap V(S_k)$ . Then, there is a  $3PC(u, v)$  unless node  $u$  has a unique neighbor  $u_i$  in  $S_i$  which is adjacent to  $v_i$  and a unique neighbor  $u_k$  in  $S_k$  which is adjacent to  $v_k$ . When this is the case, node  $u$  has an even number of neighbors in  $S_j$  (else  $(H, u)$  is an odd wheel) and  $u$  is of Type b.

Assume now that  $u$  has neighbors in exactly two sectors of the wheel, say  $S_j$  and  $S_k$ . If these two sectors are adjacent, let  $v_i$  be their common endnode and  $v_j, v_k$  the other endnodes of  $S_j$  and  $S_k$  respectively. Let  $H'$  be the hole obtained from  $H$  by replacing  $S_j \cup S_k$  by the

shortest path in  $S_j \cup S_k \cup \{u\} \setminus \{v_i\}$ . The wheel  $(H', v)$  is an odd wheel. So the sectors  $S_j$  and  $S_k$  are not adjacent. If  $u$  has three neighbors or more on  $H$ , say two or more in  $S_j$  and at least one in  $S_k$ , then denote by  $v_j$  and  $v_{j-1}$  the endnodes of  $S_j$  and by  $v_k$  one of the endnodes of  $S_k$ . There exists a  $3PC(u, v)$  where each of the nodes  $v_j, v_{j-1}$ , and  $v_k$  belongs to a different path. Therefore  $u$  has only two neighbors in  $H$ , say  $u_j \in V(S_j)$  and  $u_k \in V(S_k)$ . Let  $C_1$  and  $C_2$  be the holes formed by the node  $u$  and the two  $u_j u_k$ -subpaths of  $H$ , respectively. In order for neither  $(C_1, v)$  nor  $(C_2, v)$  to be an odd wheel, the sectors  $S_j$  and  $S_k$  must be of the same color and  $u$  is of Type a.  $\square$

*Proof of Theorem 9.5:*

Recall that  $h_1, h_3, h_5 \in V^c$  and  $h_2, h_4, h_6 \in V^r$ . We first show that  $x_1 \in V^r$  or  $x_n \in V^c$  or both. Assume the contrary, i.e.  $x_1 \in V^c$  and  $x_n \in V^r$ . Now all neighbors of  $x_1$  in  $V(\Sigma)$  are in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  and all neighbors of  $x_n$  are in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$ . Assume first that  $B(\Sigma)$  is a triad, let  $b \in V^r$  be the meet of  $B(\Sigma)$  and let  $P_2$  be the path in  $B(\Sigma) \setminus \{h_4, h_6\}$  with endnodes  $b$  and  $h_2$ .  $P_4, P_6$  are similarly defined. If  $x_n$  has neighbors in at least two of the paths, say  $P_2$  and  $P_4$ , then  $h_2, h_4$  and  $x_1$  are intermediate nodes in the three paths of a  $3PC(h_3, x_n)$ . So we can assume w.l.o.g. that all the neighbors of  $x_n$  are in  $P_2$ . Now  $h_4, h_6$  and  $x_n$  are intermediate nodes in the three paths of a  $3PC(h_5, b)$ . So, by symmetry, both  $T(\Sigma)$  and  $B(\Sigma)$  are fans. Assume  $h_6$  is the center of  $B(\Sigma)$ , so all the neighbors of  $x_n$  are in  $P_{24}$ . If  $x_n$  has more than one neighbor in  $P_{24}$ , there is a  $3PC(h_3, x_n)$ . So by symmetry  $x_1$  has a unique neighbor in  $T(\Sigma)$ , say  $x_0$  and  $x_n$  has a unique neighbor in  $B(\Sigma)$ , say  $x_{n+1}$ , but now we have a  $3PC(x_0, x_{n+1})$ . Thus we have that  $x_1 \in V^r$  or  $x_n \in V^c$ .

Since either  $x_1$  has a neighbor in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  or  $x_n$  has a neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$ , we can assume w.l.o.g. that  $x_n \in V^c$  and that  $x_1$  has a neighbor in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$ . We show that  $x_n$  has exactly two neighbors in  $\{h_2, h_4, h_6\}$  and no other neighbor in  $B(\Sigma)$ .

**Case 1:**  $B(\Sigma)$  is a triad.

Let  $b \in V^r$  be the meet of  $B(\Sigma)$ ,  $P_2$  be the path in  $B(\Sigma)$  with endnodes  $b$  and  $h_2$ .  $P_4, P_6$  are similarly defined. Let  $n_i, i = 2, 4, 6$ , be the number of neighbors of  $x_n$  in  $P_i \setminus \{b\}$ .

Assume first that  $x_n$  and  $b$  are adjacent. Then  $n_2 + n_6$  is positive, else  $h_2, h_6$  and  $x_n$  are intermediate nodes in the three paths of a  $3PC(h_1, b)$ . Now  $n_2 + n_6$  is odd, else there is an odd wheel with center  $x_n$ . By symmetry,  $n_2 + n_4$  and  $n_4 + n_6$  are also odd, but this is impossible.

Assume now that  $x_n$  and  $b$  are nonadjacent. If  $n_2, n_4$  and  $n_6$  are all positive, then we have a  $3PC(x_n, b)$ . So assume w.l.o.g.  $n_4 = 0$ . If  $n_6 = 0$ , then  $h_4, h_6$  and  $x_n$  are intermediate nodes in the three paths of a  $3PC(h_5, b)$ . So by symmetry  $n_2$  and  $n_6$  are both positive. Let  $b_2, b_6$  be respectively the neighbors of  $x_n$ , closest to  $b$  in  $P_2$  and  $P_6$ .

If  $b_2 \neq h_2$ , then  $h_3, b_2$  and  $b_6$  are intermediate nodes in the three paths of a  $3PC(x_n, b)$ . So  $b_2 = h_2$  and by symmetry,  $b_6 = h_6$  and the theorem holds in this case.

**Case 2:**  $B(\Sigma)$  is a fan.

Assume  $h_6$  is the center of the fan, let  $\ell$  be the number of neighbors of  $x_n$  in  $P_{24}$  and let  $b_1, \dots, b_k$  be the neighbors of  $h_6$ , encountered in this order when traversing  $P_{24}$  from  $h_2$  to  $h_4$ , where  $k$  is positive and even.

If  $\ell = 0$ , then  $h_6$  is the only neighbor of  $x_n$  in  $B(\Sigma)$  and  $b_1, b_k$  and  $x_n$  are intermediate nodes in the three paths of a  $3PC(h_3, h_6)$ .

If  $\ell = 1$ , let  $y_1$  be the unique neighbor of  $x_n$  in  $P_{24}$ . If  $y_1 \neq h_2$  and  $y_1 \neq h_4$ , then  $h_2$ ,  $h_4$  and  $x_n$  are intermediate nodes in the three paths of a  $3PC(h_3, y_1)$ . So we assume w.l.o.g. that  $y_1 = h_2$ . Let  $Q$  be a shortest path between  $h_5$  and  $x_n$  in  $P \cup T(\Sigma) \setminus \{h_1, h_3\}$  and let  $C = x_n, h_2, P_{24}, h_4, h_5, Q, x_n$ . Then  $x_n$  is adjacent to  $h_6$ , else  $(C, h_6)$  is an odd wheel. So  $x_n$  is adjacent to  $h_2, h_6$  and no other node in  $B(\Sigma)$ , so the theorem holds in this case.

Assume now  $\ell \geq 2$ . Then  $\ell$  is even, else  $(C, x_n)$  is an odd wheel, where  $C = h_2, P_{24}, h_4, h_3, h_2$ . Let  $y_1, y_\ell$  be the neighbors of  $x_n$ , closest to  $h_2, h_4$  in  $P_{24}$ .

Assume first that  $x_n$  is adjacent to  $h_6$ . Then  $y_1$  belongs to  $(P_{24})_{h_2b_1}$ . For, if not, let  $Q$  be a shortest path between  $h_3$  and  $x_n$ , in  $P \cup T(\Sigma) \setminus \{h_1, h_5\}$  and let  $R$  be a shortest path between  $h_1$  and  $x_n$ , in  $P \cup T(\Sigma) \setminus \{h_3, h_5\}$ . Let  $C_1 = y_1, x_n, R, h_1, h_2, (P_{24})_{h_2y_1}, y_1, x_n$ . Then  $h_6$  has an even number of neighbors in  $(P_{24})_{h_2y_1}$ , else  $(C_1, h_6)$  is an odd wheel. Let  $C_2 = y_1, x_n, Q, h_3, h_2, (P_{24})_{h_2y_1}, y_1, x_n$ . Now  $(C_2, h_6)$  is an odd wheel. So  $y_1$  belongs to  $(P_{24})_{h_2b_1}$  and by symmetry,  $y_\ell$  belongs to  $(P_{24})_{h_4b_k}$ . If  $y_1 \neq h_2$ , then the following three paths induce a  $3PC(x_n, h_2)$ :

$$x_n, y_1, (P_{24})_{y_1h_2}, h_2; \quad x_n, y_\ell, (P_{24})_{y_\ell h_4}, h_4, h_3, h_2; \quad x_n, h_6, h_1, h_2.$$

So  $y_1 = h_2$  and by symmetry,  $y_\ell = h_4$ . Let  $H = h_1, h_2, h_3, h_4, h_5, h_6, h_1$ , now  $(H, x_n)$  is an odd wheel.

Assume finally that  $\ell \geq 2$  and  $x_n$  is not adjacent to  $h_6$ . Let  $C = h_2, P_{24}, h_4, h_5, P_{51}, h_1, h_2$ . Then  $(C, h_6)$  is a wheel,  $h_6 \in V^r$  and  $x_n \in V^c$  is strongly adjacent to  $C$  and is not adjacent to  $h_6$ . First suppose that the neighbors of  $x_n$  in  $C$  are all contained in the same sector of  $(C, h_6)$ , say sector  $S$ . If  $S$  does not contain  $h_2$  nor  $h_4$ , then there is a  $3PC(x_n, h_6)$  in which two of the paths use the endnodes of  $S$  and the third path is contained in  $T(\Sigma) \cup P$ . Now w.l.o.g. assume that  $S$  contains  $h_2$ . Let  $Q$  be a shortest path between  $y_\ell$  and  $h_5$ , contained in  $P \cup T(\Sigma) \setminus \{h_1, h_3\}$  and let  $C_1 = y_\ell, Q, h_5, h_4, (P_{24})_{h_4y_\ell}, y_\ell$  then  $(C_1, h_6)$  is an odd wheel.

So no sector contains all the neighbors of  $x_n$  and Lemma 9.6 can be applied. If  $x_n$  is of Type a[9.6] with neighbors  $y_1$  and  $y_2$  in  $C$ , then since  $\ell \geq 2$ ,  $y_1, y_2 \in B(\Sigma)$ . Now  $y_1, y_2$  must coincide with  $h_2, h_4$ , else there is a  $3PC(x_n, h_6)$  and the theorem holds in this case.

So  $x_n$  is of Type b[9.6]. If all the neighbors of  $x_n$  in  $C$  belong to  $B(\Sigma)$ , there is a  $3PC(x_n, h_6)$ . If all but one of the neighbors of  $x_n$  in  $C$  belong to  $B(\Sigma)$ , then  $(C_2, x_n)$  is an odd wheel, where  $C_2 = h_3, h_2, P_{24}, h_4, h_3$ .  $\square$

**Lemma 9.7** *Let  $\Sigma$  be a connected 6-hole in a balanceable bipartite graph  $G$ . A strongly adjacent node  $w$  to  $\Sigma$  is of one of the following types:*

**Type a** *Either  $T(\Sigma)$  or  $B(\Sigma)$  contains all the neighbors of  $w$ , and  $w$  has a neighbor in  $V(\Sigma) \setminus H$ .*

**Type b** *Node  $w$  is adjacent to exactly two nodes of  $\Sigma$  and these two nodes belong to the 6-hole of  $\Sigma$ . Such a node  $w$  is called a fork.*

**Type c** *Node  $w$  has neighbors in both  $T(\Sigma)$  and  $B(\Sigma)$ , and either  $w$  has exactly two neighbors in  $\{h_1, h_3, h_5\}$  and no other neighbor in  $T(\Sigma)$ , or  $w$  has exactly two neighbors in  $\{h_2, h_4, h_6\}$  and no other neighbor in  $B(\Sigma)$ .*

*Proof:* If all the neighbors of  $w$  are either in  $T(\Sigma)$  or in  $B(\Sigma)$  and  $w$  has a neighbor in  $V(\Sigma) \setminus H$ , then  $w$  is of Type a.

If all the neighbors of  $w$  are either in  $T(\Sigma)$  or in  $B(\Sigma)$ , say in  $T(\Sigma)$  and  $w$  has no neighbor in  $V(\Sigma) \setminus H$ , then  $w$  has exactly two neighbors in  $\{h_1, h_3, h_5\}$ , else  $(H, w)$  is an odd wheel, and hence  $w$  is of Type b.

Finally, if  $w$  has neighbors in both  $T(\Sigma)$  and  $B(\Sigma)$ , then  $P = w$  is a direct connection between  $T(\Sigma)$  and  $B(\Sigma)$  in  $\Sigma \setminus E(H)$  such that  $w$  has a neighbor in  $T(\Sigma) \cup B(\Sigma) \setminus \{h_1, h_2, h_3, h_4, h_5, h_6\}$  and by Theorem 9.5,  $w$  is of Type c.  $\square$

**Lemma 9.8** *Let  $\Sigma$  be a connected 6-hole in a balanceable bipartite graph  $G$ . Every direct connection  $P = x_1, \dots, x_n$  from  $T(\Sigma)$  to  $B(\Sigma)$  in  $G \setminus E(H)$  is of one of the following types:*

- a)  $n = 1$  and  $x_1$  is a strongly adjacent node of Type c [9.7].
- b) One endnode of  $P$  is a fork, adjacent to  $h_{i-1}$  and  $h_{i+1}$ , and the other endnode of  $P$  is adjacent to a node of  $V(\Sigma) \setminus V(H)$ .
- c) Nodes  $x_1$  and  $x_n$  are not strongly adjacent to  $\Sigma$  and their unique neighbors in  $\Sigma$  are two adjacent nodes of  $H$ .
- d) One endnode of  $P$  is a fork, say  $x_1$  is adjacent to  $h_1$  and  $h_3$ , and  $x_n$  has a unique neighbor in  $\Sigma$  which is  $h_2$ .
- e) Node  $x_1$  is a fork, say adjacent to  $h_1$  and  $h_3$ , and  $x_n$  is also a fork, adjacent to  $h_2$  and either  $h_4$  or  $h_6$ .

*Proof:* If  $x_1$  or  $x_n$  has a neighbor in  $(T(\Sigma) \cup B(\Sigma)) \setminus \{h_1, h_2, h_3, h_4, h_5, h_6\}$ , by Theorem 9.5 we have a) or b). So  $n > 1$ ,  $x_1$  has no neighbor in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  and  $x_n$  has no neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$  and by Lemma 9.7,  $x_1$  and  $x_n$  are either not strongly adjacent to  $\Sigma$  or they are forks.

Assume both  $x_1$  and  $x_n$  have a unique neighbor in  $H$ , where  $x_1$  is adjacent to say  $h_1$ . If  $x_n$  is adjacent to  $h_4$  we have a  $3PC(h_1, h_4)$ , otherwise we have c).

Assume now  $x_1$  is a fork, adjacent to say  $h_1$  and  $h_3$  and  $x_n$  has a unique neighbor in  $H$ . If  $x_n$  is adjacent to  $h_2$  we have d) and if  $x_n$  is adjacent to say  $h_6$  the following three paths give a  $3PC(h_3, h_6)$ :

$$h_3, x_1, P, x_n, h_6; \quad h_3, P_{35}, h_5, h_6; \quad h_3, h_2, P_{26}, h_6.$$

Finally assume  $x_1$  is a fork, adjacent to say  $h_1$  and  $h_3$  and  $x_n$  is also a fork. If  $x_n$  is adjacent to  $h_2$  we have e). Otherwise let  $C = h_1, h_2, h_3, h_4, x_n, h_6, h_1$ . If  $n = 2$ ,  $(C, x_1)$  is an odd wheel, otherwise  $C \cup P$  contains a  $3PC(x_1, x_n)$ .  $\square$

**Lemma 9.9** *Let  $P = x_1, \dots, x_n$  be a direct connection between  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  and  $B(\Sigma)$  avoiding  $\{h_1, h_3, h_5\}$  in  $G \setminus E(H)$ , with  $x_1$  adjacent to a node in  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  and  $x_n$  adjacent to a node in  $B(\Sigma)$ . Then  $x_n$  has exactly two neighbors in  $\{h_2, h_4, h_6\}$  and no other neighbor in  $B(\Sigma)$ .*

*Proof:* Assume not and choose  $P$  and  $\Sigma$  as a counterexample to the lemma with  $P$  shortest. Now at least one intermediate node of  $P$  is adjacent to a node in  $\{h_1, h_3, h_5\}$ , else the lemma holds as a consequence of Theorem 9.5.

We show that  $x_n$  has no neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$ . Assume not and let  $x_k$  be the node of  $P$  with highest index, adjacent to a node in  $\{h_1, h_3, h_5\}$  (possibly  $k = n$ ). Then

$P_{x_k x_n}$  is a direct connection between  $T(\Sigma)$  and  $B(\Sigma)$  in  $\Sigma \setminus E(H)$  where  $x_n$  has a neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$ . So by Theorem 9.5,  $x_k$  has exactly two neighbors in  $\{h_1, h_3, h_5\}$ , say  $h_1$  and  $h_3$ , and no other neighbor in  $T(\Sigma)$ . By Lemma 9.3,  $B(\Sigma) \cup P_{x_k x_n}$  contains a tripod  $B(\Sigma')$  with attachments  $x_k, h_4$  and  $h_6$ . Let  $\Sigma'$  be a connected 6-hole with top  $T(\Sigma') = T(\Sigma)$  and bottom  $B(\Sigma')$ . ( $\Sigma'$  exists by Lemma 9.4). Now  $P_{x_1 x_{k-1}}$  is a direct connection from  $T(\Sigma') \setminus \{h_1, h_3, h_5\}$  to  $B(\Sigma')$  avoiding  $\{h_1, h_3, h_5\}$  and, since  $x_k$  is the unique neighbor of  $x_{k-1}$  in  $B(\Sigma')$ , this contradicts our choice of  $P$  and  $\Sigma$ .

So we assume w.l.o.g. that node  $x_n$  is adjacent to  $h_2$  and no other node of  $B(\Sigma)$ .

We now show that at most two of the nodes in  $\{h_1, h_3, h_5\}$  have neighbors in  $P$ . If all three nodes  $h_1, h_3, h_5$  have neighbors in  $P$ , then by Lemma 9.3, there exists a tripod  $T(\Sigma')$  contained in  $(P \setminus \{x_n\}) \cup \{h_1, h_3, h_5\}$  with attachments  $h_1, h_3$  and  $h_5$ . Let  $\Sigma'$  be a connected 6-hole with top  $T(\Sigma')$  and bottom  $B(\Sigma') = B(\Sigma)$  (Again,  $\Sigma'$  exists by Lemma 9.4). Now a subpath of  $P$  is a direct connection between  $T(\Sigma') \setminus \{h_1, h_3, h_5\}$  and  $B(\Sigma')$  avoiding  $\{h_1, h_3, h_5\}$  and this contradicts our choice of  $P$  and  $\Sigma$ .

We show that  $h_5$  has no neighbors in  $P$ . Assume not and let  $x_l$  be the node of highest index adjacent to  $h_5$ . By the previous argument, either  $h_1$  or  $h_3$  has no neighbors in  $P$ . W.l.o.g. assume node  $h_3$  has no neighbors in  $P$ . Now there exists a  $3PC(h_5, h_2)$  with paths  $h_5, x_l, P_{x_l x_n}, x_n, h_2$ ;  $h_5, P_{53}, h_3, h_2$  and  $h_5, h_6, P_{62}, h_2$ .

So we assume w.l.o.g. that  $h_1$  is adjacent to an intermediate node of  $P$  while  $h_5$  is not. Let  $Q$  be a shortest path in  $\{x_1\} \cup T(\Sigma) \setminus \{h_1, h_3\}$  between  $x_1$  and  $h_5$ . Now one of the two holes  $x_n, P, x_1, Q, h_5, h_6, P_{62}, x_n$  or  $x_n, P, x_1, Q, h_5, h_4, P_{42}, x_n$  induces an odd wheel with center  $h_1$ .  $\square$

## 9.2 Extreme Connected 6-Holes

**Definition 9.10** *An extreme connected 6-hole  $E\Sigma$  is a subgraph of  $G$  containing six nonempty node sets  $H_1, \dots, H_6$  such that, if  $H$  is the graph induced by  $H_1 \cup \dots \cup H_6$ , then  $E(H)$  is a 6-join of  $E\Sigma$  separating subgraphs  $T$  and  $B$ , where  $V(T) \cup V(B) = V(E\Sigma)$ ,  $H_1 \cup H_3 \cup H_5 \subset V(T)$ ,  $H_2 \cup H_4 \cup H_6 \subset V(B)$  and the three following properties hold:*

- (1) *Let  $T'_1, \dots, T'_m$  be the connected components of the graph  $T'$  induced by  $V(T) \setminus (H_1 \cup H_3 \cup H_5)$ . Then  $m \geq 1$ , each  $T'_j$  has at least one neighbor in each of the sets  $H_1, H_3, H_5$  and each node in  $H_1 \cup H_3 \cup H_5$  has at least one neighbor in  $T'$ .*

*The graph  $B'$  induced by  $V(B) \setminus (H_2 \cup H_4 \cup H_6)$  is nonempty and connected. Each node in  $H_2 \cup H_4 \cup H_6$  has at least one neighbor in  $B'$ .*

- (2) *For  $i = 1, 3, 5$  and  $j = 1, \dots, m$ , let  $H_i^j$  be the set of nodes in  $H_i$  with a neighbor in  $T'_j$  and let  $T_j$  be the graph induced by the node set  $V(T'_j) \cup H_1^j \cup H_3^j \cup H_5^j$ . Let the  $H$ -intersection graph of  $E\Sigma$  be defined as follows: its node set is  $\{t_1, \dots, t_m\}$  and  $t_j$  is adjacent to  $t_k$  if at least two of the following three sets are nonempty:*

$$H_1^j \cap H_1^k, \quad H_3^j \cap H_3^k, \quad H_5^j \cap H_5^k.$$

*Then the  $H$ -intersection graph of  $E\Sigma$  is connected.*

(3)  $V(E\Sigma)$  is maximal, subject to (1) and (2).

**Lemma 9.11** *The graphs  $T_j$  and  $B$  satisfy the following properties:*

(1) *For every index  $j$  and triple of nodes  $h_1 \in H_1^j, h_3 \in H_3^j, h_5 \in H_5^j$ ,  $T_j$  contains a fan or a triad with attachments  $h_1, h_3, h_5$  and all other nodes in  $T_j'$ .*

*For every triple of nodes  $h_2 \in H_2, h_4 \in H_4, h_6 \in H_6$ ,  $B$  contains a fan or a triad with attachments  $h_2, h_4, h_6$  and all other nodes in  $B'$ .*

(2) *Let  $S$  be a fan or a triad in  $T_j$  satisfying (1) and  $s$  be any node of  $T_j \setminus S$ . Then either  $s$  is adjacent to a node in  $S \setminus \{h_1, h_3, h_5\}$  or  $T_j'$  contains a direct connection between  $s$  and  $S \setminus \{h_1, h_3, h_5\}$ .*

*Let  $R$  be a fan or a triad in  $B$  satisfying (1) and  $r$  be any node of  $B \setminus R$ . Then either  $r$  is adjacent to a node in  $R \setminus \{h_2, h_4, h_6\}$  or  $B'$  contains a direct connection between  $r$  and  $R \setminus \{h_2, h_4, h_6\}$ .*

*Proof:* By definition,  $T_j'$  is a connected graph and since  $h_1, h_3, h_5$  all have neighbors in  $T_j'$ , then the graph induced by  $V(T_j') \cup \{h_1, h_3, h_5\}$  is also connected. So by Lemma 9.3,  $T_j$  contains a tripod  $S$  with attachments  $h_1, h_3, h_5$  and all other nodes in  $V(T_j')$ . The same argument shows that for every three nodes  $h_2 \in H_2, h_4 \in H_4, h_6 \in H_6$ ,  $B$  contains a tripod  $R$  with attachments  $h_2, h_4, h_6$  and all other nodes in  $B'$ . Now by Lemma 9.4 applied to the graph induced by  $V(S) \cup V(R)$ , we have that  $S$  and  $R$  are indeed fans or triads and (1) follows.

Since  $S \setminus \{h_1, h_3, h_5\} \subseteq T_j'$  and by definition  $V(T_j') \cup \{s\}$  induces a connected graph, then the first part of (2) follows. The proof of the second part is identical.  $\square$

**Theorem 9.12** *Let  $E\Sigma$  be an extreme connected 6-hole in a balanceable bipartite graph  $G$  and let  $U$  be a connected component of  $G \setminus V(E\Sigma)$ , with neighbors in  $T$  and in  $B$ . Then  $U$  has no neighbor in  $T' \cup B'$ .*

*Proof:* Assume not. Since  $U$  has neighbors of both  $T$  and  $B$ , then either  $U$  contains a direct connection between  $T'$  and  $B$  avoiding  $H_1 \cup H_3 \cup H_5$ , or  $U$  contains a direct connection between  $T$  and  $B'$  avoiding  $H_2 \cup H_4 \cup H_6$  (or both). Among all these direct connections, let  $Q = y_1, \dots, y_\ell$  be a shortest one. (Possibly  $\ell = 1$ ).

**Case 1:** Node  $y_1$  has a neighbor in  $T'$ ,  $y_\ell$  has a neighbor in  $B$  and no intermediate node of  $Q$  is adjacent to a node in  $H_2 \cup H_4 \cup H_6$ .

We assume that  $y_1$  has a neighbor in  $T_j'$ .

**Claim 1:** Node  $y_\ell$  has no neighbor in  $B'$ .

*Proof of Claim 1:* Assume  $y_\ell$  has a neighbor in  $B'$ . Let  $T(\Sigma)$  be any fan or triad in  $T_j$  with attachments  $h_1 \in H_1^j, h_3 \in H_3^j, h_5 \in H_5^j$  and  $B(\Sigma)$  be any fan or triad in  $B$  with attachments  $h_2 \in H_2, h_4 \in H_4, h_6 \in H_6$ . Let  $\Sigma$  be the connected 6-hole in  $E\Sigma$  with  $T(\Sigma)$  as top and  $B(\Sigma)$  as bottom. (By Lemma 9.11(1) such a  $\Sigma$  exists). By Lemma 9.11(2), there exists a direct connection  $P = x_1, \dots, x_n$  from  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  to  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$  avoiding  $\{h_1, h_2, h_3, h_4, h_5, h_6\}$ , such that  $P = P_{T_j'}, Q, P_{B'}$  where  $P_{T_j'} \subset T_j'$  and  $P_{B'} \subset B'$ . Possibly  $P_{T_j'}$

or  $P_{B'}$  or both are empty. If no intermediate node of  $P$  is adjacent to a node in  $\{h_2, h_4, h_6\}$ , then  $P$  is a direct connection from  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  and  $B(\Sigma)$ , avoiding  $\{h_1, h_3, h_5\}$  and, since  $x_n$  has a neighbor in  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$ ,  $P$  contradicts Lemma 9.9. So at least one intermediate node of  $P$  has a neighbor in  $\{h_2, h_4, h_6\}$  and the same argument shows that at least one intermediate node of  $P$  has a neighbor in  $\{h_1, h_3, h_5\}$ . Let  $x_r$  be the intermediate node of  $P$  with highest index with a neighbor in  $\{h_1, h_3, h_5\}$ , and let  $x_s$  be the intermediate node of  $P$  with lowest index with a neighbor in  $\{h_2, h_4, h_6\}$ . By construction, the intermediate nodes of  $P$  that have neighbors in  $\{h_1, h_3, h_5\}$  belong to  $P_{T'_j}$  or  $Q$  and the intermediate nodes of  $P$  that have neighbors in  $\{h_2, h_4, h_6\}$  are either  $y_\ell$  or belong to  $P_{B'}$ . Clearly  $y_\ell$  cannot have neighbors in both  $\{h_1, h_3, h_5\}$  and  $\{h_2, h_4, h_6\}$ . This shows that  $r < s$ . Now  $P_{x_1 x_s}$  is a direct connection from  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  to  $B(\Sigma)$ , avoiding  $\{h_1, h_3, h_5\}$  and by Lemma 9.9,  $x_s$  has exactly two neighbors in  $B(\Sigma)$ , say  $h_2$  and  $h_6$ . By Lemma 9.3,  $T_j \cup P_{x_1 x_s}$  contains a tripod  $T(\Sigma')$  with attachments  $x_s, h_3, h_5$  and all other nodes in  $T'_j \cup P_{x_1 x_{s-1}}$ . Let  $\Sigma'$  be the connected 6-hole with top  $T(\Sigma')$  and bottom  $B(\Sigma') = B(\Sigma)$ . Now  $P_{x_{s+1} x_n}$  is a direct connection from  $T(\Sigma')$  to  $B(\Sigma') \setminus \{h_2, h_4, h_6\}$  avoiding  $\{h_2, h_4, h_6\}$  and  $x_s$  is the unique neighbor of  $x_{s+1}$  in  $T(\Sigma')$ , a contradiction to Lemma 9.9. This completes the proof of Claim 1.

**Claim 2:** Node  $y_\ell$  is adjacent to all nodes in exactly two of the sets  $H_2, H_4, H_6$  and to no other node of  $B$ .

*Proof of Claim 2:* By Claim 1,  $N(y_\ell) \cap V(B) \subseteq H_2 \cup H_4 \cup H_6$ . Assume  $y_\ell$  is adjacent to  $h_2 \in H_2$ . Let  $B(\Sigma)$  be any fan or triad in  $B$  with attachments  $h_2, h_4 \in H_4, h_6 \in H_6$ , let  $T(\Sigma)$  be any fan or triad in  $T_j$  having attachments  $h_1 \in H_1^j, h_3 \in H_3^j, h_5 \in H_5^j$  and let  $\Sigma$  be the connected 6-hole with top  $T(\Sigma)$  and bottom  $B(\Sigma)$ . Such a choice of  $\Sigma$  is possible by Lemma 9.11(1). Now by Lemma 9.11(2), there exists a direct connection  $R$  from  $T(\Sigma) \setminus \{h_1, h_3, h_5\}$  to  $B(\Sigma)$  avoiding  $\{h_1, h_3, h_5\}$ , such that  $R = R_{T'_j}, Q$ , where  $V(R_{T'_j}) \subset V(T'_j)$  and possibly  $V(R_{T'_j})$  is empty. So by Lemma 9.9 applied to  $\Sigma$  and  $R$ ,  $y_\ell$  has exactly two neighbors in  $B(\Sigma)$ , say  $h_2$  and  $h_6$ .

Let  $\Sigma'$  be any connected 6-hole with top  $T(\Sigma') = T(\Sigma)$  and bottom  $B(\Sigma')$  with exactly two common attachments with  $B(\Sigma)$ . (Again by Lemma 9.11(1),  $\Sigma'$  exists.) Now  $R$  is a direct connection from  $T(\Sigma') \setminus \{h_1, h_3, h_5\}$  to  $B(\Sigma')$  avoiding  $\{h_1, h_3, h_5\}$ . So by Lemma 9.9,  $y_\ell$  is adjacent to the new attachment  $h'$  of  $B(\Sigma')$  if and only if  $h' \in H_2 \cup H_6$ . By Lemma 9.11(1), every node  $h'$  in  $H_2 \cup H_4 \cup H_6 \setminus \{h_2, h_4, h_6\}$  is the attachment of such  $B(\Sigma')$ . So, by Lemma 9.9,  $y_\ell$  is adjacent to all nodes in  $H_2 \cup H_6$  and to no other node of  $B$  and this completes the proof of Claim 2.

By Claim 2 we may assume w.l.o.g. that  $y_\ell$  is adjacent to all nodes in  $H_2 \cup H_6$  and to no other node of  $B$ . Let  $E\Sigma^*$  be the subgraph of  $G$ , induced by  $V(E\Sigma) \cup V(Q)$ , where  $H_1^* = H_1 \cup \{y_\ell\}$ ,  $H_i^* = H_i$  for all the other indices, and let  $H^*$  the graph induced by  $H_1^* \cup \dots \cup H_6^*$ . By Claim 2,  $E(H^*)$  is a 6-join of  $E\Sigma^*$ , separating  $T^* = T \cup Q$  from  $B^* = B$ .

If  $\ell = 1$ , then let  $T_k^{k*} = T'_k$  for  $k = 1, \dots, m$ , let  $H_1^{k*} = H_1^k \cup \{y_1\}$  if and only if  $y_1$  has a neighbor in  $T'_k$  and  $H_i^{k*} = H_i^k$  in all other cases. By construction,  $T_1^{k*}, \dots, T_m^{k*}$  are the connected components of the graph induced by  $V(T') \setminus (H_1^* \cup H_3^* \cup H_5^*)$  and  $H_i^{k*}$  contains the nodes in  $H_i^*$  with a neighbor in  $T_k^{k*}$ . The  $H^*$ -intersection graph of  $E\Sigma^*$  is connected since  $H_i^k \subseteq H_i^{k*}$  for all  $k$  and  $i = 1, 3, 5$ . So  $E\Sigma^*$  satisfies Properties (1) and (2) of Definition 9.10, contradicting the assumption that  $E\Sigma$  is extreme.

If  $\ell > 1$ , assume w.l.o.g. that  $y_1$  has no neighbor in  $T'_1, \dots, T'_{p-1}$  and has at least one neighbor in each of  $T'_p, \dots, T'_m$ . Let  $T_1^{p*} = T'_1, \dots, T_{p-1}^{p*} = T'_{p-1}$  and let  $T_p^{p*}$  be the connected component induced by  $V(T'_p) \cup \dots \cup V(T'_m) \cup \{y_1, \dots, y_{\ell-1}\}$ . For  $k = 1, \dots, p-1$ , let  $H_i^{k*} = H_i^k$ . Finally, let  $H_1^{p*}$  contain  $\{y_\ell\} \cup \cup_{k=p}^m H_1^k$  together with all nodes of  $H_1$  with a neighbor in  $Q_{y_1 y_{\ell-1}}$ , and for  $i = 3, 5$ , let  $H_i^{p*}$  contain  $\cup_{k=p}^m H_i^k$  together with all nodes in  $H_i$  with a neighbor in  $Q_{y_1 y_{\ell-1}}$ . By construction,  $T_1^{p*}, \dots, T_p^{p*}$  are the connected components of the graph induced by  $V(T^*) \setminus (H_1^* \cup H_3^* \cup H_5^*)$  and  $H_i^{k*}$  contains the nodes in  $H_i^*$  with a neighbor in  $T_k^{p*}$ .

The  $H^*$ -intersection graph of  $E\Sigma^*$  is connected since for  $i = 1, 3, 5$ ,  $H_i^k \subseteq H_i^{k*}$  for all  $k = 1, \dots, p-1$ , and  $H_i^k \subseteq H_i^{p*}$  for all  $k = p, \dots, m$ . From this it follows that  $E\Sigma^*$  satisfies Properties (1) and (2) of Definition 9.10, a contradiction to the assumption that  $E\Sigma$  is extreme.

**Case 2:** Node  $y_1$  has a neighbor in  $T$ ,  $y_\ell$  has a neighbor in  $B'$  and no intermediate node of  $Q$  is adjacent to a node in  $H_1 \cup H_3 \cup H_5$ .

The same proof given for Claim 1 shows that  $y_1$  has no neighbor in  $T'$ , so  $N(y_1) \cap V(T) \subseteq H_1 \cup H_3 \cup H_5$ .

**Claim 3:** Node  $y_1$  is adjacent to all the nodes in exactly two of the sets  $H_1, H_3, H_5$  and to no other node of  $T$ .

*Proof of Claim 3:* By Case 1 we may assume that  $y_1$  is not adjacent to a node of  $T'$ . W.l.o.g. assume  $y_1$  has a neighbor in  $h_1 \in H_1^j$ , let  $T(\Sigma)$  be any fan or triad in  $T_j$  having attachments  $h_1, h_3 \in H_3^j, h_5 \in H_5^j, B(\Sigma)$  be any fan or triad in  $B$  with attachments  $h_2 \in H_2, h_4 \in H_4, h_6 \in H_6$  and let  $\Sigma$  be a connected 6-hole with top  $T(\Sigma)$  and bottom  $B(\Sigma)$ . Now by Lemma 9.11(2), there exists a direct connection  $R$  from  $T(\Sigma)$  and  $B(\Sigma) \setminus \{h_2, h_4, h_6\}$  avoiding  $\{h_2, h_4, h_6\}$ , such that  $R = Q, R_{B'}$ , where  $V(R_{B'}) \subset V(B')$  and possibly  $V(R_{B'})$  is empty. So by Lemma 9.9 applied to  $\Sigma$  and  $R$ ,  $y_1$  has exactly two neighbors in  $T(\Sigma)$ , say  $h_1$  and  $h_3$ . Now the same argument used in the proof of Claim 2 shows that  $y_1$  is adjacent to all the nodes in  $H_1^j \cup H_3^j$  and no other node of  $T_j$ .

Choose now  $T_k$  such that at least two of the following three sets are nonempty:  $H_1^j \cap H_1^k, H_3^j \cap H_3^k, H_5^j \cap H_5^k$ . (This choice is possible by Property (2) of Definition 9.10). Let  $T(\Sigma')$  be any fan or triad in  $T_k$  having attachments  $h'_1 \in H_1^k, h'_3 \in H_3^k, h'_5 \in H_5^k$ , where at least two of these attachments are in  $T_j$  and let  $\Sigma'$  be a connected 6-hole with top  $T(\Sigma')$  and bottom  $B(\Sigma') = B(\Sigma)$ . (Lemma 9.11(1) shows that  $\Sigma'$  exists). Now since at least two of the attachments of  $T(\Sigma')$  are in  $T_j$  and  $y_1$  is adjacent to all the nodes in  $H_1^j \cup H_3^j$  and no other node of  $T_j$ , by Lemma 9.9 applied to  $\Sigma'$  and  $R$ , we have that  $h'_1$  and  $h'_3$  are the unique neighbors of  $y_1$  in  $T(\Sigma')$ . This shows that  $y_1$  is adjacent to all the nodes in  $H_1^k \cup H_3^k$  and no other node of  $T_k$ . Now by Property (2) of Definition 9.10, we obtain that  $y_1$  is adjacent to all the nodes in  $H_1 \cup H_3$  and no other node of  $T$  and the proof of Claim 3 is complete.

Let  $E\Sigma^*$  be the subgraph of  $G$ , induced by  $V(E\Sigma) \cup V(Q)$ , where  $H_2^* = H_2 \cup \{y_1\}$ ,  $H_i^* = H_i$  for all the other indices and  $H^*$  is the subgraph induced by  $H_1^* \cup \dots \cup H_6^*$ . By Claim 3,  $E(H^*)$  is a 6-join of  $E\Sigma^*$ , separating  $T^* = T$  from  $B^* = B \cup Q$ . Now  $V(B^*) \setminus (H_2^* \cup H_4^* \cup H_6^*)$  induces a connected graph, since  $B'$  is connected and  $y_\ell$  has a neighbor in  $B'$ . So  $E\Sigma^*$  satisfies Properties (1) and (2) of Definition 9.10, a contradiction to the fact that  $E\Sigma$  is extreme.  $\square$

### 9.3 Extended Star Cutsets and 6-Joins

**Lemma 9.13** *Let  $\Sigma$  be a connected 6-hole in a balanceable bipartite graph  $G$  and let  $i \in \{1, 2, 3\}$ . Let  $P$  be a direct connection from  $h_i$  to  $h_{i+3}$  such that the nodes of  $P$  are in  $G \setminus V(\Sigma)$ , have no neighbors in  $V(\Sigma) \setminus \{h_1, \dots, h_6\}$  and no proper subpath of  $P$  is a direct connection from  $h_j$  to  $h_{j+3}$ , for  $j \in \{1, 2, 3\}$ . Then exactly two of the nodes  $h_{i-2}, h_{i-1}, h_{i+1}, h_{i+2}$ , have a neighbor in  $P$  and these two nodes are either  $\{h_{i+1}, h_{i+2}\}$  or  $\{h_{i-1}, h_{i-2}\}$ .*

*Proof:* Let  $P = x_1, \dots, x_n$  with  $x_1$  adjacent to  $h_i$  and  $x_n$  to  $h_{i+3}$ . Assume w.l.o.g. that  $i = 3$ .

We first show that if  $P$  contains no neighbor of  $h_5$ , then  $P$  contains neighbors of  $h_2$  and  $h_1$ .

If  $P$  contains no neighbors of  $h_2$ , there exists a  $3PC(h_3, h_6)$  where the three paths are  $P$ ;  $h_3, P_{35}, h_6$  and  $h_3, h_2, P_{26}, h_6$ . So  $P$  must contain a neighbor of  $h_2$ . If  $P$  contains no neighbors of  $h_1$ , then one of the two holes  $h_3, P, h_6, h_1, P_{13}, h_3$  or  $h_3, P, h_6, h_5, P_{53}, h_3$  makes an odd wheel with center  $h_2$ .

We now show that if  $P$  contains no neighbor of  $h_5$ , then  $P$  contains no neighbor of  $h_4$ .

Since  $P$  contains no neighbor of  $h_5$ , then  $P$  contains neighbors of  $h_2$  and  $h_1$ . If  $P$  has a neighbor of  $h_4$ , then there exists a direct connection  $P'$  from  $h_1$  to  $h_4$  in  $G \setminus V(\Sigma)$  using nodes in  $P$ . By minimality of  $P$ ,  $P = P'$ . Thus  $x_1$  is adjacent to  $h_1$  and  $x_n$  to  $h_4$ , and nodes  $h_1$  and  $h_4$  have no other neighbors in  $P$ . Let  $x_j$  be the neighbor of  $h_2$  with the highest index. Then  $x_j, \dots, x_n, h_4, h_5, P_{51}, h_1, h_2, x_j$  makes an odd wheel with center  $h_6$ .

So, if node  $h_5$  has no neighbors in  $V(P)$ , the lemma holds. Now by symmetry, if any one of the nodes  $\{h_1, h_2, h_4, h_5\}$  has no neighbors in  $P$ , we are done. If all four nodes have neighbors in  $P$ , then  $P$  contains a direct connection  $P'$  from  $h_1$  to  $h_4$  in  $G \setminus V(\Sigma)$  and a direct connection  $P''$  from  $h_5$  to  $h_2$  in  $G \setminus V(\Sigma)$ . By minimality of  $P$ ,  $P = P' = P''$ . But then  $x_1$  is adjacent to  $h_1, h_3$  and  $h_5$ . Consequently  $(H, x_1)$  is an odd wheel.  $\square$

**Theorem 9.14** *Let  $E\Sigma$  be an extreme connected 6-hole in a balanceable bipartite graph  $G$  and let  $U$  be a connected component of  $G \setminus V(E\Sigma)$ , with neighbors in both  $T$  and in  $B$ . If for some  $i$ , both  $H_i$  and  $H_{i+3}$  contain neighbors of  $U$ , then there exists an extended star cutset, separating at least one node of  $U$  from  $E\Sigma$ .*

*Proof:* Let  $U$  be a connected component of  $G \setminus V(E\Sigma)$  with neighbors in  $H_i$  and  $H_{i+3}$  for some  $i = 1, 2$  or  $3$ . By Theorem 9.12, all the neighbors of  $U$  in  $E\Sigma$  belong to  $H$ . So  $U$  contains a direct connection from  $H_i$  and  $H_{i+3}$  with no neighbor in  $V(E\Sigma) \setminus H$ . Among all these direct connections and possible choices of  $i$ , let  $P = x_1, \dots, x_n$  be a shortest one and assume w.l.o.g. that  $x_1$  is adjacent to a node  $h_3 \in H_3^j$  and  $x_n$  to a node  $h_6 \in H_6$ .

**Claim 1:** Either every node in  $H_1^j \cup H_2$  has a neighbor in  $P$  and no node in  $H_4 \cup H_5$  has a neighbor in  $P$ , or every node in  $H_4 \cup H_5^j$  has a neighbor in  $P$  and no node in  $H_1 \cup H_2$  has a neighbor in  $P$ .

*Proof of Claim 1:* For every pair of nodes  $h_1 \in H_1^j$  and  $h_5 \in H_5^j$ , let  $T(\Sigma)$  be a fan or a triad in  $T$ ; with attachments  $h_1, h_3, h_5$ . For every pair of nodes  $h_2 \in H_2$  and  $h_4 \in H_4$ , let  $B(\Sigma)$  be a fan or a triad in  $B$  with attachments  $h_2, h_4, h_6$  and let  $\Sigma$  be the connected 6-hole with  $T(\Sigma)$  as top and  $B(\Sigma)$  as bottom. By Lemma 9.13, we can assume that both  $h_1$  and  $h_2$  have neighbors in  $P$ , while  $h_4$  and  $h_5$  do not have neighbors in  $P$ . By Lemma 9.11(1), it

follows readily that every node in  $H_1^j \cup H_2$  has a neighbor in  $P$  and no node in  $H_3^j \cup H_4$  has a neighbor in  $P$ .

It remains to show that no node in  $H_5 \setminus H_5^j$  is adjacent to a node of  $P$ . Assume not and let  $h'_5 \in H_5^k \setminus H_5^j$  be such a node. Let  $\Sigma'$  be a connected 6-hole having top  $T(\Sigma') \subset T_k$  with attachments  $h'_5$  and arbitrarily chosen nodes  $h'_1 \in H_1^k$  and  $h'_3 \in H_3^k$  and bottom  $B(\Sigma') = B(\Sigma)$ . By the previous argument,  $h_2$  has a neighbor in  $P$ , while  $h_4$  has no neighbor in  $P$ . So  $P$  contains a direct connection  $P'$  between  $h'_5$  and  $h_2$  and by the minimality of  $P$ ,  $P' = P$ . So  $x_1$  is the unique neighbor of  $h'_5$  in  $P$  and  $x_n$  is the unique neighbor of  $h_2$  in  $P$ . Now, by Lemma 9.13 applied to  $\Sigma'$ ,  $P$  and  $i = 2$ , node  $h'_1$  has a neighbor in  $P$ , since  $h_4$  has no neighbor in  $P$ . Let  $x_j$  be a neighbor of  $h'_1$  with lowest index. Since  $x_n$  is the unique neighbor of  $h_2$  or  $h_6$  in  $P$ , then  $P_{x_1 x_j}$  contains no neighbor in  $\{h_2, h_4, h_6\}$ . Let  $H^*$  be the 6-hole  $h'_1, h_2, h_3, h_4, h'_5, h_6$ , and consider the graph  $G^*$  induced by  $V(H^*) \cup V(P_{x_1 x_j})$ . Then  $x_1$  is adjacent to  $h'_5, h_3$  and  $x_j$  to  $h'_1$ . So if  $j = 1$ ,  $G^*$  is an odd wheel with center  $x_1$  and, if  $j > 1$ ,  $G^*$  contains a  $3PC(x_1, h'_1)$  and this completes the proof of Claim 1.

By Claim 1, we can assume w.l.o.g. that every node in  $H_1^j \cup H_2$  has a neighbor in  $P$  and no node in  $H_4 \cup H_5$  has a neighbor in  $P$ . Let  $h_2^*$  be a node in  $H_2$ . Let  $S$  be the extended star  $(h_2^*; H_2; H_1 \cup H_3; N(h_2^*) \setminus V(E\Sigma))$ . We show that  $S$  is an extended star cutset separating  $x_1$  from  $E\Sigma$ .

Assume not. Then the connected component  $U$  contains a direct connection  $Q = y_1, \dots, y_q$  from  $x_1$  to  $V(E\Sigma) \setminus V(S)$ , avoiding  $V(S)$ . Since  $y_q$  belongs to  $U$ , by Theorem 9.12,  $N(y_q) \cap V(E\Sigma) \subset V(H)$ . Let  $Q' = x_1, y_1, \dots, y_q$ .

**Case 1:**  $y_q$  is adjacent to a node  $h'_6 \in H_6$ .

Let  $\Sigma$  be a connected 6-hole containing  $h_2^*, h_3, h'_6$  and arbitrary other attachments  $h_1 \in H_1^j, h_4 \in H_4, h_5 \in H_5^j$ . Let  $Q''$  be a minimal subpath of  $Q'$  which is a direct connection from  $h_3$  to  $h'_6$  or from  $h_1$  to  $h_4$ . Since  $Q''$  has no neighbors in  $V(\Sigma) \setminus \{h_1, h_3, h_4, h'_6\}$ , it is a minimal direct connection satisfying the assumptions of Lemma 9.13 relative to  $\Sigma$ . But then, by Lemma 9.13,  $h_2^*$  or  $h_5$  has a neighbor in  $Q''$ , contradicting the choice of  $Q$  or the assumption that  $x_1$  is not adjacent to  $h_5$ .

**Case 2:**  $y_q$  is adjacent to a node  $h_4 \in H_4$ .

We can assume that  $y_q$  is not adjacent to any node in  $H_6$ . Therefore no node in  $H_5 \cup H_6$  has a neighbor in  $Q'$ . Let  $R = r_1(= y_q), \dots, r_t$  be a direct connection from  $h_4$  to  $H_1^j$  with  $V(R) \subseteq V(P) \cup V(Q)$  (such  $R$  exists since  $P$  has at least one neighbor in  $H_1^j$ ). We can assume w.l.o.g. that  $r_1, \dots, r_s \in V(Q) \setminus V(P)$  for some  $s \leq t$  and  $r_j \in V(P)$  for  $j > s$ . Let  $h_1 \in H_1^j$  be a neighbor of  $r_t$  and let  $\Sigma$  be a connected 6-hole with attachments  $h_1, h_2^*, h_3, h_4, h_6$  and an arbitrary node  $h_5 \in H_5^j$ . Node  $h_5$  has no neighbor in  $R$ . Furthermore,  $h_6$  has no neighbor in  $R$  since  $x_n$  is the only node of  $V(P) \cup V(Q)$  adjacent to  $h_6$  and  $R$  cannot contain  $x_n$ . Therefore  $R$  is a minimal set of nodes satisfying the assumptions of Lemma 9.13 relative to  $\Sigma$ . This implies that both  $h_2^*$  and  $h_3$  have neighbors in  $R$ . Let  $C_1 = h_4, r_1, R, r_t, h_1, h_6, P_{64}, h_4$  ( $P_{64}$  is defined in Remark 9.1). Then  $R$  has an odd number of neighbors of  $h_3$ , else  $(C_1, h_3)$  is an odd wheel. No node of  $Q'$  is adjacent to  $h_1$  since, otherwise, some subpath of  $Q'$  would be a direct connection from  $h_1$  to  $h_4$  violating Lemma 9.13 in  $\Sigma$  (since neither  $h_2^*$  nor  $h_6$  has a neighbor in  $Q'$ ). Therefore, by construction of  $R$ , if  $r_\ell$  denotes the node of lowest index adjacent to  $h_2^*$ , then all the neighbors of  $h_3$  in  $R$  are in  $R_{r_1 r_\ell}$ . Let  $C_2 = h_4, r_1, R_{r_1 r_\ell}, r_\ell, h_2^*, h_1, P_{15}, h_5, h_4$ .

Since  $h_3$  has an even number of neighbors in  $P_{15}$  and an odd number of neighbors in  $R_{r_1 r_\ell}$ , then  $(C_2, h_3)$  is an odd wheel.

**Case 3:**  $y_q$  is adjacent to a node  $h_5 \in H_5^k$ .

Then  $Q$  has no neighbors of  $H_4 \cup H_6$ . We first show that no node in  $H_2$  is adjacent to a node of  $Q$ . Let  $y_s$  be the node of highest index adjacent to a node  $t_2 \in H_2$ . Let  $\Sigma$  be a connected 6-hole containing  $t_2, h_5, h_6$  and arbitrary other attachments  $h_1 \in H_1^k, h'_3 \in H_3^k, h_4 \in H_4$ . Now  $Q_{y_s y_q} \subseteq U$  is a subpath of  $Q$ , and is a direct connection from  $t_2$  to  $h_5$  satisfying the assumptions of Lemma 9.13 relative to  $\Sigma$ . So  $Q_{y_s y_q}$  must contain a neighbor of  $h_4$  or  $h_6$ , which is a contradiction. So  $Q$  has no neighbor in  $H_2 \cup H_4 \cup H_6$ .

Let  $x_\ell$  be the node of  $P$  with lowest index adjacent to a node in  $H_2$ , say  $t_2$ . Let  $\Sigma$  be a connected 6-hole with attachments  $t_2, h_5, h_6$  and arbitrary  $h_1 \in H_1^k, h'_3 \in H_3^k, h_4 \in H_4$ . Now  $P_{x_1 x_\ell} \cup Q$  contains a direct connection  $P'$  from  $t_2$  to  $h_5$  in  $G \setminus V(\Sigma)$  with no neighbors in  $V(\Sigma) \setminus \{h_1, t_2, h'_3, h_4, h_5, h_6\}$ . Either  $P'$  is a direct connection from  $t_2$  to  $h_5$  satisfying the assumptions of Lemma 9.13 relative to  $\Sigma$ , or a subpath  $P''$  of  $P'$  is a direct connection from  $h'_3$  to  $h_6$  satisfying these assumptions (these are the only two possibilities since  $h_4$  has no neighbor in  $P'$ ). In both cases, Lemma 9.13 implies that  $P'$  contains a neighbor of  $h_1$  and that  $x_\ell = x_n$  (since  $h_4$  has no neighbor in  $P'$  and  $h_6$  is only adjacent to  $x_n$  in  $P'$  or  $P''$ ). But now the nodes of  $P_{x_1 x_{n-1}} \cup Q$  have no neighbors in  $H_2$ . So the nodes of  $P_{x_1 x_{n-1}} \cup Q$  have no neighbors in  $B$ . Since the graph induced by  $V(P_{x_1 x_{n-1}}) \cup V(Q) \cup \{h_1, h_3, h_5\}$  is connected, by Lemma 9.3, there exists a tripod  $Y$  with attachments  $h_1, h_3$  and  $h_5$ , contained in  $V(P_{x_1 x_{n-1}}) \cup V(Q) \cup \{h_1, h_3, h_5\}$ .

Let  $E\Sigma^*$  be the subgraph of  $G$ , induced by  $V(E\Sigma) \cup V(Y)$ . Let  $H_i^* = H_i$  and let  $H^*$  be the graph induced by  $H_1^* \cup \dots \cup H_6^*$ . Then  $E(H^*)$  is a 6-join of  $E\Sigma^*$ , separating  $T^* = T \cup Y$  from  $B^* = B$ . Now the connected components of  $V(T^*) \setminus \{H_1^* \cup H_3^* \cup H_5^*\}$  are the same as the ones for  $E\Sigma^*$  except for a new one, namely  $Y^* = Y \setminus \{h_1, h_3, h_5\}$ . Let  $H_i^Y$  denote the set of neighbors of  $Y^*$  in  $H_i$ , for  $i = 1, 3, 5$ . Since  $h_1 \in H_1^Y \cap H_1^k$  and  $h_5 \in H_5^Y \cap H_5^k$ , the  $H^*$ -intersection graph of  $E\Sigma^*$  is connected. It follows that  $E\Sigma^*$  satisfies Properties (1) and (2) of Definition 9.10, a contradiction to the fact that  $E\Sigma$  is extreme.  $\square$

Now we can prove Theorem 6.4.

**Theorem 6.4** *A balanceable bipartite graph  $G$  that contains a connected 6-hole as an induced subgraph, has an extended star cutset or a 6-join.*

*Proof:* Since a connected 6-hole satisfies (1) and (2) of Definition 9.10, the assumption that  $G$  contains a connected 6-hole implies that  $G$  contains an extreme connected 6-hole  $E\Sigma$ . Let  $U_1, \dots, U_k$  be the connected components of  $G \setminus V(E\Sigma)$  having at least one neighbor in  $T$  and at least one neighbor in  $B$ . Note that  $E(H)$  is a 6-join of  $G$ , separating  $T$  and  $B$  and only if no such component exists. By Theorem 9.12, no connected component  $U_j$  has a neighbor in  $T' \cup B'$ , so  $H$  contains all the neighbors of  $U_j$ . If all the neighbors of  $U_j$  belong to  $H_{i-1} \cup H_i \cup H_{i+1}$  for some  $i$ , then  $K_{H_i, H_{i-1} \cup H_{i+1}}$  is a biclique cutset, separating  $U_j$  and  $E\Sigma$ . Otherwise  $U_j$  has neighbors in  $H_i$  and in  $H_{i+3}$ , for some  $i$ . Now, by Theorem 9.14, there exists an extended star cutset, separating at least one node of  $U_j$  from  $E\Sigma$ .  $\square$

## References

- [1] C. Berge, Sur certains hypergraphes généralisant les graphes bipartites, in: *Combinatorial Theory and its Applications I* (P. Erdős, A. Rényi and V. Sós eds.), *Colloq. Math. Soc. János Bolyai 4*, North Holland, Amsterdam (1970) 119-133.
- [2] C. Berge, Balanced matrices, *Mathematical Programming 2* (1972) 19-31.
- [3] P. Camion, Characterization of totally unimodular matrices, *Proceedings of the American Mathematical Society 16* (1965) 1068-1073.
- [4] V. Chandru and J.N. Hooker, Extended Horn sets in propositional logic, *Journal of the ACM 38* (1991) 205-221.
- [5] M. Conforti and G. Cornuéjols, A class of logic problems solvable by linear programming, *Journal of the ACM 42* (1995) 1107-1113.
- [6] M. Conforti and G. Cornuéjols, Balanced 0,  $\pm 1$  matrices, bicoloring and total dual integrality, *Mathematical Programming 71* (1995) 249-258.
- [7] M. Conforti, G. Cornuéjols, A. Kapoor, M. R. Rao, K. Vušković, Balanced matrices, in *Mathematical Programming: State of the Art 1994*, J.R. Birge and K.G. Murty eds., The University of Michigan Press (1994) 1-33.
- [8] M. Conforti, G. Cornuéjols and M. R. Rao, Decomposition of balanced matrices, *Journal of Combinatorial Theory B 77* (1999) 292-406.
- [9] M. Conforti and M. R. Rao, Structural properties and recognition of restricted and strongly unimodular matrices, *Mathematical Programming 38* (1987) 17-27.
- [10] G. Cornuéjols and W. H. Cunningham, Compositions for perfect graphs, *Discrete Mathematics 55* (1985) 245-254.
- [11] W. H. Cunningham and J. Edmonds, A combinatorial decomposition theory, *Canadian Journal of Mathematics 32* (1980) 734-765.
- [12] D. R. Fulkerson, A. Hoffman and R. Oppenheim, On balanced matrices, *Mathematical Programming Study 1* (1974) 120-132.
- [13] G. Georgakopoulos, D. Kavvadias and C. H. Papadimitriou, Probabilistic satisfiability, *Journal of Complexity 4* (1988) 1-11.
- [14] A. Ghouila-Houri, Caractérisations des matrices totalement unimodulaires, *C. R. Acad. Sc. Paris 254* (1962) 1192-1193.
- [15] P. Seymour, Decomposition of regular matroids, *Journal of Combinatorial Theory B 28* (1980) 305-359.
- [16] K. Truemper, Alpha-balanced graphs and matrices and GF(3)-representability of matroids, *Journal of Combinatorial Theory B 32* (1982) 112-139.

- [17] K. Truemper, Polynomial theorem proving I. Central matrices, *Technical Report UTDCS 34-90* (1990).
- [18] K. Truemper, *Matroid Decomposition*, Academic Press, Boston (1992).